

EXPRESS MAIL NO. EV 320 858 167 US

DATE OF MAILING 2/7/03

Attorney Docket No. 9800080-0054-Cont

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE
APPLICATION FOR UNITED STATES LETTERS PATENT

INVENTORS:

David P. Andrew
David A. Lewin
Diane Pennica
Luca Rastelli
Bruce Tallion

TITLE:

WNT-Regulated Cytokine-like Polypeptide
and Nucleic Acids Encoding Same

ATTORNEY

Paul E. Rauch, Ph.D.
SONNENSCHN NATH & ROSENTHAL LLP
P. O. Box 061080
Wacker Drive Station
Sears Tower
Chicago, Illinois 60606
(312) 876-8000

WNT-REGULATED CYTOKINE-LIKE POLYPEPTIDE AND NUCLEIC ACIDS ENCODING SAME

RELATED APPLICATIONS

This application claims priority to USSN __, filed November 16, 2000 and USSN
5 60/166,177, filed November 17, 1999. The contents of these applications are incorporated by
reference in their entireties.

FIELD OF THE INVENTION

The invention relates in general to polynucleotides and polypeptides. The invention
10 relates more particularly to polynucleotide sequences and the membrane-bound or secreted
polypeptides encoded by such polynucleotides, as well as vectors, host cells, antibodies and
recombinant methods for producing the polypeptides and polynucleotides.

BACKGROUND OF THE INVENTION

15 Wnt proteins are a family of secreted polypeptides that act through a signaling
pathway, which is known as the Wnt signaling pathway. Previously described members of the
Wnt signalling pathway include the putative Wnt receptor, which is encoded by members of
the frizzled gene family. Other members of the Wnt-signalling pathway include the
transcription factor beta-catenin, the adenomatous polyposis coli (APC) polypeptide,
20 presenilin (PS), GSK-3b, Protein Phosphatase 2A proteins, the kinases CamKII and PKC,
glycogen synthase kinase 3, the Dishevelled (Dvl) protein, and cyclins D1 and c-myc.
Transcription of these latter genes is reported to be associated with activation of the Wnt
signaling pathway. Interactions between some members of the Wnt-signaling pathway have
been described. For example, APC is phosphorylated by GSK-3b, binds to beta-catenin and
25 facilitates its degradation.

Wnt proteins control cell processes via at least two pathways. One of these, known as
the "canonical" Wnt signaling pathway, operates through the cytosolic stabilization of beta-
catenin (Smalley MJ and Dale TC, 1999). Stabilization is achieved by downregulating the
activity of a beta-catenin turnover complex. In the second pathway, activation of the Wnt
30 signaling pathway stimulates intracellular Ca²⁺ release and activates the kinases CamKII and
PKC (Kuhl et al, Trends Genet 2000 Jul;16(7):279-83).

Wnt signaling is involved in a variety of mammalian developmental processes,
including cell proliferation, differentiation and epithelial-mesenchymal interactions (Smalley

MJ and Dale TC, Cancer Metastasis Rev 1999;18(2):215-30). For example, Wnt signaling contributes to the development of tissues and organs such as the limbs, the brain, the reproductive tract and the kidney during embryogenesis (Peifer M and Polakis P, Science 2000 Mar 3;287(5458):1606-9).

5 Constituents of Wnt signaling pathways are expressed in embryogenesis and act in developing and mature nervous systems (Patapoutian A, and Reichardt LF, Curr Opin Neurobiol 2000 Jun;10(3):392-9). For example, Wnt signaling is involved in the initial formation of the neural plate and in subsequent patterning decisions in the embryonic nervous system, including formation of the neural crest. Wnt signaling is also required at later stages of
10 development.

 Actions of members of the Wnt signaling pathway have also been associated with disease processes. For example, Wnt signaling has been demonstrated to regulate apoptosis and may participate in degenerative processes leading to cell death in the aging brain.

 In addition, inappropriate activation of the canonical Wnt signaling pathway is
15 reported to be a major feature in human neoplasia. Oncogenic activation of this pathway can occur in several ways. For example, mutations in either APC or beta-catenin have been associated with colon carcinomas. Inappropriate expression of Wnt and Wnt binding proteins has similarly been found in a variety of human tumors. Dysregulation of the beta-catenin turnover complex has been associated with several tumor types. Thus dysregulation can
20 include in, *e.g.*, loss of the Adenomatous Polyposis Coli (APC) or Protein Phosphatase 2A proteins, or activating mutations of beta-catenin.

 Disruption of expression of members of the Wnt signaling pathway has also been associated with apoptosis and neurological conditions such as Alzheimer's disease (De Ferrari GV, and Inestrosa NC, Brain Res Brain Res Rev 2000 Aug;33(1):1-12).

25

SUMMARY OF THE INVENTION

 The present invention is based in part upon the discovery of a nucleic acid sequence encoding a novel member of the Wnt signaling pathway. The novel member encodes a S100 cytokine-like polypeptide. The S100 cytokine-like nucleic acids and polypeptides encoded
30 thereby are collectively referred to herein as "FCRX".

 Accordingly, in one aspect, the invention provides an isolated nucleic acid molecule that encodes a novel polypeptide, or a fragment, homolog, analog or derivative thereof. The nucleic acid can include, *e.g.*, a nucleic acid sequence encoding a polypeptide at least 85% identical to a polypeptide comprising the amino acid sequences of SEQ ID NO: 3 or 6, or a

polypeptide that is a fragment, homolog, analog or derivative thereof. The nucleic acid can include, *e.g.*, one or more fragments from genomic DNA, or a cDNA molecule, or an RNA molecule. In particular embodiments, the nucleic acid molecule may include the sequence of any of SEQ ID NO: 1, 2, 4, or 5. These polypeptides and nucleic acids are related to S100
5 cytokine-like proteins, as disclosed herein.

Also included in the invention is a vector containing one or more of the nucleic acids described herein, and a cell containing the vectors or nucleic acids described herein.

The invention is also directed to host cells transformed with a vector comprising any of the nucleic acid molecules described above.

10 In another aspect, the invention includes a pharmaceutical composition that includes an FCTR_X nucleic acid and a pharmaceutically acceptable carrier or diluent.

In a further aspect, the invention includes a substantially purified FCTR_X polypeptide, *e.g.*, any of the FCTR_X polypeptides encoded by an FCTR_X nucleic acid, and fragments, homologs, analogs, and derivatives thereof. The invention also includes a pharmaceutical
15 composition that includes an FCTR_X polypeptide and a pharmaceutically acceptable carrier or diluent.

In a still further aspect, the invention provides an antibody that binds specifically to an FCTR_X polypeptide. The antibody can be, *e.g.*, a monoclonal or polyclonal antibody, and fragments, homologs, analogs, and derivatives thereof. The invention also includes a
20 pharmaceutical composition including FCTR_X antibody and a pharmaceutically acceptable carrier or diluent. The invention is also directed to isolated antibodies that bind to an epitope on a polypeptide encoded by any of the nucleic acid molecules described above. The invention also includes kits comprising any of the pharmaceutical compositions described above.

25 The invention further provides a method for producing an FCTR_X polypeptide by providing a cell containing an FCTR_X nucleic acid, *e.g.*, a vector that includes an FCTR_X nucleic acid, and culturing the cell under conditions sufficient to express the FCTR_X polypeptide encoded by the nucleic acid. The expressed FCTR_X polypeptide is then recovered from the cell. Preferably, the cell produces little or no endogenous FCTR_X
30 polypeptide. The cell can be, *e.g.*, a prokaryotic cell or eukaryotic cell.

The invention is also directed to methods of identifying a FCTR_X polypeptides or nucleic acids in a sample by contacting the sample with a compound that specifically binds to the polypeptide or nucleic acid, and detecting complex formation, if present.

The invention further provides methods of identifying a compound that modulates the activity of an FCTR_X polypeptide by contacting FCTR_X polypeptide with a compound and determining whether the FCTR_X polypeptide activity is modified.

5 The invention is also directed to compounds that modulate FCTR_X polypeptide activity identified by contacting an FCTR_X polypeptide with the compound and determining whether the compound modifies activity of the FCTR_X polypeptide, binds to the FCTR_X polypeptide, or binds to a nucleic acid molecule encoding an FCTR_X polypeptide.

10 In another aspect, the invention provides a method of determining the presence of or predisposition to an FCTR_X-associated disorder in a subject. The method includes providing a sample from the subject and measuring the amount of FCTR_X polypeptide in the subject sample. The amount of FCTR_X polypeptide in the subject sample is then compared to the amount of FCTR_X polypeptide in a control sample. An alteration in the amount of FCTR_X polypeptide in the subject protein sample relative to the amount of FCTR_X polypeptide in the control protein sample indicates the subject has pathology related to a dysfunction in the
15 immune system, a tissue proliferation-associated condition, or a neurological disorder. A control sample is preferably taken from a matched individual, *i.e.*, an individual of similar age, sex, or other general condition but who is not suspected of having a dysfunction in the immune system, a tissue proliferation-associated condition, or a neurological disorder. Alternatively, the control sample may be taken from the subject at a time when the subject is not suspected
20 of having a dysfunction in the immune system, a tissue proliferation-associated condition, or a neurological disorder. In some embodiments, the FCTR_X polypeptide is detected using an FCTR_X antibody.

In a further aspect, the invention provides a method of determining the presence of, or predisposition to an FCTR_X-associated disorder in a subject. The method includes providing
25 a nucleic acid sample, *e.g.*, RNA or DNA, or both, from the subject and measuring the amount of the FCTR_X nucleic acid in the subject nucleic acid sample. The amount of FCTR_X nucleic acid sample in the subject nucleic acid is then compared to the amount of FCTR_X nucleic acid in a control sample. An alteration in the amount of FCTR_X nucleic acid in the sample relative to the amount of FCTR_X in the control sample indicates the subject has a dysfunction in the
30 immune system, a tissue proliferation-associated condition, or a neurological disorder.

In a still further aspect, the invention provides a method of treating or preventing or delaying an FCTR_X-associated disorder. The method includes administering to a subject in which such treatment or prevention or delay is desired an FCTR_X nucleic acid, an FCTR_X polypeptide, or an FCTR_X antibody in an amount sufficient to treat, prevent, or delay an

immune disorder, a tissue proliferation-associated disorder, or a neurological disorder in the subject.

Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, suitable methods and materials are described below. All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety. In the case of conflict, the present specification, including definitions, will control. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

Other features and advantages of the invention will be apparent from the following detailed description and claims.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a ClustalW alignment of the human polypeptide of SEQ ID NO:6 with known S100 cytokine proteins.

Figure 2 is a ClustalW alignment of the human polypeptide of SEQ ID NO:6 and the murine ortholog of SEQ ID NO:3.

Figure 3 is a Pfam analysis of human EST AA315020 (SEQ ID NO:4).

Figure 4A is a BLOCKS protein domain analysis of the polypeptide of SEQ ID NO:6.

Figure 4B shows an alignment of amino acids 28-131 of SEQ ID NO:3 with amino acids 1-101 of Acc. No. AY007220, an S-100 type calcium binding protein.

Figure 4C shows an alignment of amino acids 28-131 of SEQ ID NO:6 with amino acids 1-101 of Acc. No. AY007220.

Figure 4D shows an alignment of a region of the polypeptide of SEQ ID NO:3 with various calcium binding domains.

Figure 4E shows an alignment of a region of the polypeptide of SEQ ID NO:6 with various previously described calcium binding domains.

Figure 5 is a hydrophobicity plot of the polypeptide of SEQ ID NO:6.

Figure 6 is a representation of the differential gene expression in transgenic Wnt-1 mice (Set A) compared to wild type mice (Set B) detected by labeled fragments run in capillary electrophoresis.

Figure 7 is a CAP alignment and consensus of human assembly 65677221-3-frag (SEQ ID NO:5) compared to human EST AA315020 (SEQ ID NO:4).

DETAILED DESCRIPTION OF THE INVENTION

The present invention discloses genes in the Wnt signaling pathway. Included in the invention is a nucleic acid expressed at different levels in murine mammary tumors that arise spontaneously in Wnt-1 transgenic mice relative to control tissues. The sequence of this differentially expressed sequence is shown in Table 1 (SEQ ID NO:1).

Table 1. Nucleotide sequence of murine clone 7971c.7.

GAATTCAGTGATGTAGAGAGGGCCATTGAGACACTCATCAAGAACTTCCATAAATACTCTGTG
GCGGGTAAAAAGGAAACACTGACCCCTGCTGAGCTTCGAGACCTGGTTACCCAGCAGCTGCCA
CACCTCATGCCGAGCAACTGTGGGTTAGAAGAGAAAATTGCCAACCTGGGCAACTGTAATGAC
TCGAAACTGGAGTTTGGGAAGCTT (SEQ ID NO:1)

The sequence shown in Table 1 was extended by assembly with other murine nucleic acid fragments. The resulting novel assembly, designated 7971c.7-r0s0-212.2-EXT (SEQ ID NO:2), is shown in Table 2.

Table 2. Nucleotide sequence of murine clone 7971c.7-r0s0-212.2-EXT.

TCAGGTGAGCTGGCTCCTCCATCCTGTCTCCCAGCTGCCAGCAGGTCTCCCCCTCCTCTAGGT
AGATCATGATCCATCAGCTCCTGTGGGGCAGGCTATAGGACAGACGACAAAACCTCAACTCACA
GAAGGAAGGACCAGTGTACCAGGAACGATGGGACAGTGTGGTCAGCCAATGCTGAGGATGCC
CAAGAATTCAGTGATGTAGAGAGGGCCATTGAGACACTCATCAAGAACTTCCATAAATACTCT
GTGGCGGGTAAAAAGGAAACACTGACCCCTGCTGAGCTTCGAGACCTGGTTACCCAGCAGCTG
CCACACCTCATGCCGAGCAACTGTGGGTTAGAAGAGAAAATTGCCAACCTGGGCAACTGTAAT

GACTCGAAACTGGAGTTTGGGAAGCTTCTGGGAGTTGATTGGAGAAGCAGCCAAGAGTGTGAAG
 ATGGAGAGGCCTGTTACTCGGAGCTGAGGACTTCTACTTGGAAGTTGTTGGGGGTGTTGGGGA
 TAGGGGAGTTTTAGAGGCACTGGAAATAAAACCTCAATGCCACCACCCCTTCCCCAGCCT
 GCACCTCTCCTCATTGCTGCAATGTTACGTTGAGGACAGGCTTCCCTGTGGGCTCCATGGAG
 5 CTCCTGGGTCCAGAAGTCCTCATCTCAAGGGAGCTCAGGGGGTGGGTGTTGGGGCTGGAGAGGAT
 ATGCAGGGATCCTGGAAGGGTAAGGGCCAAGCAATTTGGTAGTAGGGGAAGGGCAGAAAGGAA
 CTGGGTTATGGAAGTGATCCAAAGAGCAGGGATGGGAATCTGGCTGCATATTTGGTCCTGAAA
 AGGGTGTCTGAGAACCTACCCCTTCTAATCTTGTCCCACCTAACTGTAGTTGTCTGCCCTG
 TGCTATCCTTGCTGCTTCCAGCTCTGCCCCATCCTCCTTCCAGTGTCTGTTCTGAGTAGGGG
 10 CAGGGGAAATAGGAGCAGAGTTGCAAAGAGGCTGAGGAGGGCATGACTTCATCACTTTGGGG
 TGAGAGGACCAGCTAGATGCTTGGGCATTTATGGTAGTTATTTTATATCATTTGATTAATAAA
 AATATTGGAAAATGTAAAGAAAAAAAAGAAAAAACATGGGGCCGAAACCTTATCCCCCTTG
 AGTAGGGTGATATTTTGCCTGTGCAATGGGCGGCCTGTTTTCGAGAGGCGGTGACATGGGGAA
 AACATGGGGGTGTACCAAACCTTAACCGCCTTTTAGGGGAAACACCCCTTTTGCCGCAAGTGG
 15 GTTAATAACGGAAGAAGCCCGCGGATTGCCCTTACAAGAGTCTCCCGCGGTAGATGCGGA
 TGGGACAGCCCCCTTCGGCGGCGTTTAGAGCGGCGTGTGTGTGGTTTCTACGCGAATAGGGAT
 AAATATTGTGGCGGCGCCGAGGGAGTGTGTGTGTTGCGCGCTGCTTCTGTGGAGGTGGTGTG
 TCCCCAAAATAAAAGGGCCCTTTTGTGCGCGTTAGTTTGCTCTAGCAGAGTCCGCTGCACAT
 ATTTTGGTGGGCGTGTCCGTGCCGCCCCGNGGTGGTGCTTGTTGCTGGCGTGGCGTGGGGTGGG
 20 TGTGGTTGCGGGGGTGGTCGTGTTGGGTGTGTGCGTGCGCGGGGGCCGTGTGTGTGTGTGG
 TTGCATGATAAGGTTAGAGTGAGTGAGAGCGG (SEQ ID NO:2)

The murine sequence shown in Table 2 includes an open reading frame ("ORF") from
 nucleotide 73 to nucleotide 465. The beginning of this ORF is indicated in bold in Table 2.

25 The polypeptide encoded by this ORF is shown in Table 3 (SEQ ID NO:3).

Table 3. Translation product of murine clone 7971c.7-r0s0-212.2-EXT.

SISSCGAGYRTDDKTQLTEGRTSVPGTMGQCRSANAEDAQEFSDVERAIETLIKFNHKEYSVAG
 KKETLTPAELRDLVTQQLPHLMPSNCGLEEKIANLGNCSKLEFGSFWELIGEAAKSVKMER
 30 PVTRS (SEQ ID NO:3)

The murine sequence shown in Table 1 is related to human EST AA315020. This EST
 originates from human cells forming a metastatic tumor when implanted in mice. The
 sequence of this EST is shown in Table 4 (SEQ ID NO:4).

35

Table 4. Nucleotide sequence of human EST AA315020.

ATAGGACAACAGAACTCTCACCAAAGGACCAGACACAGTGAGCACCATGGGACAGTGTCTGGTC
AGCCAACGCAGAGGATGCTCAGGAATTCAGTGATGTGGAGAGGGCCATTGAGACCCTCATCAA
GAACTTTCACCAGTACTCCGTGGAGGGTGGGAAGGAGACGCTGACCCCTTCTGAGCTACGGGA
5 CCTGGTCACCCAGCAGCTGCCCCATCTCATGCCGAGCAACTNTGGCCTGGAAGAGAAAATTGC
CAACCTGGGCAGCTGCAATGACTCTAACTGGAGTTCAGGAGTTTCTGGGAGCTGATTGGAGA
AGCGGCCAAGAGTGTGAAGCTNGAGAGGACTGTCCGGGGGCA (SEQ ID NO:4)

A consensus sequence (SEQ ID NO:5) of 294 bp was assembled as described in
10 Example 2. The consensus sequence shown in Table 5 encompasses and extends human EST
AA315020 (SEQ ID NO:4).

Table 5. Human Consensus Sequence 65677221-3-frag

GAATTCCAGAGGGAGTTCTCAGTGCCCCCGGACAGGCCTCTCCAGCTTCACACTCTTGGCCGC
15 TTCTCCAATCAGCTCCCAGAACTCCTGAACTCCAGTTTAGAGTCATTGCAGCTGCCAGGTT
GGCAATTTTCTCTCCAGGCCACAGTTGCTCGGCATGAGATGGGGCAGCTGCTGGGTGACCAG
GTCCCGTAGCTCAGAAGGGGTCAGCGTCTCCTTCCCACCCTCCACGGAGTACTGGTGAAAGTT
CTTGATGAGGGTCTCAATGGCCCTCTCCACATCACTGAATTCCTGAGCATCCTCTGCGTTGGC
TGACCGACACTGTCCCATGGTGCTCACTGTGTCTGGTCCTTTGGTGAGAGTTCTGTTGTCCTA
20 T (SEQ ID NO:5)

The nucleotide sequence shown in Table 5 includes an ORF that encodes a translation
product having the sequence shown in Table 6 (SEQ ID NO:6).

25 Table 6. Predicted translation of 65677221-3-frag.

DNRTLTKGPDVSTMGQCRSANAEDAQEFSDVERAIETLIKNFHQYSVEGGKETLTPSELRDL
VTQQLPHLMPSNCGLEEKIANLGSCNDSKLEFRSFWELIGEAAKSVKLERPVRGH (SEQ ID
NO:6)

The polypeptides whose sequences are shown in Table 3 and Table 6 have a high
30 degree of similarity with the S100 family of proteins. FIG. 1 demonstrates this similarity for
the human sequence shown in Table 6. The sequence identified as "W27152" in Fig. 1 is
disclosed as a "chemotactic cytokine II CCII polypeptide" in PCT publication WO97/34013,
and the polypeptide designated "G491246" is a
Macrophage Migration Inhibition Factor (MRP-14).

The mouse and human amino acid sequences of Table 3 and Table 6 are shown aligned in Fig. 2. This alignment reveals that these two sequences are sufficiently similar such that they can be considered to be orthologs. The region of similarity between the human and the mouse proteins begins just before this first Met of the human sequence and ends just before the stop codon. In addition, the codon encoding the first Met is located near a Kozak consensus sequence. Analysis by the Pfam algorithm that identifies related protein families indicates that the polypeptide encoded by the human EST AA315020 (SEQ ID NO:4) is related to the S100 family (Fig. 3).

Moreover, analysis by the BLOCKS program (see Fig. 4A) shows the two conserved calcium binding regions separated by about 8 amino acids, which is a characteristic of S100 proteins (Kligman. 1988 Trends Biochem. Sci. 13:437).

Figs.4B-4E show comparisons of portions of the amino acid sequences shown in Table 3 (SEQ ID NO:3) and Table 6 (SEQ ID NO:6) to S-100-type calcium binding proteins. Fig. 4B shows an alignment of amino acids 28-131 of SEQ ID NO:3 with amino acids 1-101 of Acc. No. AY007220, a S-100 type calcium binding protein. Identical or conserved amino acid residues are indicated in black shading. These residues may be required to preserve structural or functional properties of the protein. Amino acids shaded in gray can be mutated to a residue with comparable steric and/or chemical properties without altering protein structure or function, *e.g.*, L to V, I or M. Non-highlighted amino acid residues can potentially be mutated to a much broader extent without altering structure or function.

Fig. 4C shows an alignment of amino acids 28-131 of SEQ ID NO:6 with amino acids 1-101 of Acc. No. AY007220. Shading is as explained for FIG. 4B.

Fig. 4B and 4C demonstrate that the polypeptides shown in Table 3 and 6 include extensive regions identical or conserved with the amino acid sequence of Acc. No. AY007220.

The polypeptide sequences shown in Tables 3 and 6 include sequences highly homologous to previously-described S-100/ICaBP-type calcium binding domains. These calcium binding domains are discussed in Sastry et al., Structure 6:223-31, 1998. Proteins containing these domains are implicated in cell growth and division. In addition, expression of these proteins is reported to be deregulated in transformed cells. For example, high levels of these proteins have been reported in some breast cancers.

Fig. 4D shows an alignment of a region of the polypeptide of Table 3 with various calcium binding domains. Fig. 4E shows an alignment of a region of the polypeptide of Table 6 with various previously described calcium binding domains.

The hydrophobicity plot of this polypeptide, shown in Fig.5, suggests that its amino acid sequence is highly hydrophilic, so that it is disposed neither toward membrane localization nor toward secretion. This is confirmed by PSORT analysis, which predicts that the polypeptide localizes in the cytoplasm. Cytoplasmic localization is another characteristic of S100 proteins (Dale et al. 1983 Eur. J. Biochem. 134:1-6). S100 proteins fall into a category of proteins that have extracellular function but are released, rather than being secreted.

It is expected that the serum concentration of a S100 cytokine according to the invention (*e.g.*, a polypeptide having the amino acid sequence of SEQ ID NO:3 or SEQ ID NO:6) can be used as a marker for the clinical predictive value for metastatic tumors. Notably, the mouse gene fragment disclosed in Table 1 (SEQ ID NO:1) was originally identified in the tumors of Wnt-1 transgenic mice. In addition, two of the human fragments used to produce the sequence shown in Table 5 (SEQ ID NO:5) were isolated from tumors. In addition, as is shown in Example 3, the sequence shown in Table 5 (SEQ ID NO:5) is highly expressed in several tumor cell lines, especially in colon cancer, breast cancer and ovarian cancer. For all these reasons, the nucleic acids and their gene products identified herein are useful as diagnostic markers for such cancers, and as targets for the treatment of cancer in a subject. The targeting may especially be carried out by generating antibodies that specifically bind a protein encoded by SEQ ID NO:5, and administering such antibodies to a subject suffering from a cancer such as colon cancer, breast cancer or ovarian cancer. These antibodies will likewise serve as diagnostic tools useful in detecting the presence or amount of the proteins disclosed herein.

A FCTR_X polypeptide according to the invention includes a polypeptide including the amino acid sequences of SEQ ID NO:3 or SEQ ID NO:6, as well as homologs, and variants as described herein. A FCTR_X nucleic acid according to the invention includes the nucleic acids shown in SEQ ID NO:1, 2, 4, or 5 as well as homologs and variants disclosed herein.

S100 Cytokine Polypeptides

The FCTR_X polypeptides are homologous to members of the S100 cytokine family. The S100 group of proteins are calcium-binding molecules with cytokine and chemokine activity. For example, the murine protein S100A8 is a potent chemoattractant for neutrophils and monocytes *in vivo* and *in vitro* (Xu K and Geczy CL, J Immunol 2000 May 1;164(9):4916-23).

The S100 family of Ca²⁺-binding proteins that contain EF-hand Ca²⁺ binding motifs includes 19 members that are differentially expressed in a large number of cell types. Members of this protein family have been implicated in the Ca²⁺-dependent regulation of

intracellular activities as diverse as protein phosphorylation, enzyme activities, cell proliferation (including neoplastic transformation) and differentiation, cytoskeleton and membrane remodeling, the structural organization of membranes, intracellular Ca^{2+} homeostasis, inflammation, and in protection from oxidative cell damage. S100 proteins have also been implicated in Zn^{2+} - or Cu^{2+} -dependent activities. Some S100 members are released or secreted into the extracellular space and exert trophic or toxic effects (depending on their concentration). Family members can also act as chemoattractants for leukocytes, modulate cell proliferation, or regulate macrophage activation (Donato. 1999 Biochim Biophys Acta 8:191-231). The serum concentration of S100 has been shown to have clinical predictive value for metastatic tumors.

S100 family members have also been reported to interact with each other. For example, recombinantly expressed murine S100A9 has been reported to interact *in vitro* with murine and human S100A8 (Nacken et al., Eur J Biochem 2000 Jan;267(2):560-5). Both murine S100A9 and its dimerization partner mS100A8 are expressed at the onset of granulocyte-colony stimulating factor induced myeloid differentiation. Substantial amounts of this complex are constitutively secreted by granulocytic 32D cells into the medium. It has also been suggested that the human and murine S100A9 may share a higher degree of functional homology than of sequence similarity (Nacken et al., 2000).

Two S100 family proteins, S100B and S100A1, activate RAGE (the receptor for an advanced glycation end product (AGE)) occurring in diabetes in concert with amphoterin. This interaction is reported to induce neurite outgrowth and activation of transcription factor NF-kappaB. Amphoterin is a protein that enhances process extension and migration in embryonic neurons and in tumor cells through binding to RAGE, a multiligand transmembrane receptor. Furthermore, activation of RAGE by amphoterin and S100B promotes cell survival through increased expression of the anti-apoptotic protein Bcl-2. Higher concentrations of S100B induce apoptosis in an oxidant-dependent manner. Activation of RAGE by multiple ligands may promote trophic effects, whereas hyperactivation of RAGE signaling pathways promotes apoptosis.

Specific staining of WHO grade II and III astrocytic tumors with S100 calcium-binding proteins has been used to characterize distinct clinical entities in both WHO grade II and III astrocytic tumors. The grade II and III astrocytic tumors lead to different clinical outcomes (Camby et al., Neuropathol Appl Neurobiol 2000 Feb;26(1):76-90). The level of cell proliferation (determined by means of both anti-proliferating cell nuclear antigen and anti-MIB-1 antibodies differed significantly between astrocytomas and anaplastic astrocytomas. In

sharp contrast, the levels of expression of the S100A3 and S100A5 proteins differed markedly in the solid tumor tissue in relation to the astrocytic tumor types and grades. In addition, while the levels of expression of S100A6 did not change in the astrocytic tumor tissue in relation to histopathological grade, the levels of expression of S100A6 (but not those of S100A3 and S100A5) differed markedly in the blood vessel walls according to whether these vessels originated from low- or high-grade astrocytic tumors. Thus S100 proteins have highly specific expression levels and degrees of association with varying cell types and tissues.

Noncancerous CNS pathologies are also related to S100 proteins. Homozygous APPV717F transgenic mice overexpress a human beta-amyloid precursor protein (betaAPP) minigene encoding a familial Alzheimer's disease mutation. These mice develop Alzheimer-type neuritic beta-amyloid plaques surrounded by astrocytes. S100beta is an astrocyte-derived cytokine that promotes neurite growth and promotes excessive expression of betaAPP. S100beta overexpression in Alzheimer's disease correlates with the proliferation of betaAPP-immunoreactive neurites in beta-amyloid plaques. Age-related increases in tissue levels of both betaAPP and S100beta mRNA have been reported in transgenic mice (Sheng et al., J Neurochem 2000 Jan;74(1):295-301). Accelerated age-related overexpression of S100beta may interact with age-associated overexpression of mutant betaAPP in transgenic mice to promote development of Alzheimer-like neuropathological changes.

Levels of S100 beta in serum is also a significant prognostic marker for malignant melanoma (Djukanovic etl al., Anticancer Res 2000 May-Jun;20(3B):2203-7). In 53 melanoma patients a direct correlation of S100 values and clinical course could be observed (81.5%). S100 beta levels were incorrectly elevated in 5 out of 25 sera, giving "false positive" results (20%) and in 8 out of 40 sera were not elevated despite the detection of metastases giving "false negative" results (20%).

FCTR X Nucleic Acids

One aspect of the invention pertains to isolated nucleic acid molecules that encode FCTR X proteins or biologically active portions thereof, as well as nucleic acid fragments sufficient for use as hybridization probes to identify FCTR X-encoding nucleic acids (*e.g.*, FCTR X mRNA) and fragments for use as PCR primers for the amplification or mutation of FCTR X nucleic acid molecules.

As used herein, the term "nucleic acid molecule" is intended to include DNA molecules (*e.g.*, cDNA or genomic DNA), RNA molecules (*e.g.*, mRNA), analogs of the DNA or RNA generated using nucleotide analogs, and derivatives, fragments and homologs thereof.

The nucleic acid molecule can be single-stranded or double-stranded, but preferably is double-stranded DNA.

An "isolated" nucleic acid molecule is one that is separated from other nucleic acid molecules which are present in the natural source of the nucleic acid. Preferably, an "isolated" nucleic acid is free of sequences which naturally flank the nucleic acid (*i.e.*, sequences located at the 5' and 3' ends of the nucleic acid) in the genomic DNA of the organism from which the nucleic acid is derived. For example, in various embodiments, the isolated FCTR_X nucleic acid molecule can contain less than about 5 kb, 4 kb, 3 kb, 2 kb, 1 kb, 0.5 kb or 0.1 kb of nucleotide sequences which naturally flank the nucleic acid molecule in genomic DNA of the cell from which the nucleic acid is derived. Moreover, an "isolated" nucleic acid molecule, such as a cDNA molecule, can be substantially free of other cellular material or culture medium when produced by recombinant techniques, or of chemical precursors or other chemicals when chemically synthesized.

In some embodiments, the FCTR_X nucleic acids encode a mature form a FCTR_X polypeptide. As used herein, the term a "mature" form of a polypeptide or protein is the product of a naturally occurring polypeptide or precursor form or FCTR_X-protein. The naturally occurring polypeptide, precursor or FCTR_X-protein includes, by way of non-limiting example, the full length gene product, encoded by the corresponding gene. Alternatively, it may be defined as the polypeptide, precursor or FCTR_X-protein encoded by an open reading frame described herein. The product "mature" form arises, again by way of non-limiting example, as a result of one or more naturally occurring processing steps as they may take place within the cell, or host cell, in which the gene product arises. Examples of such processing steps leading to a "mature" form of a polypeptide or protein include the cleavage of the N-terminal methionine residue encoded by the initiation codon of an open reading frame, or the proteolytic cleavage of a signal peptide or leader sequence. Thus a mature form arising from a precursor polypeptide or protein that has residues 1 to N, where residue 1 is the N-terminal methionine, would have residues 2 through N remaining after removal of the N-terminal methionine. Alternatively, a mature form arising from a precursor polypeptide or protein having residues 1 to N, in which an N-terminal signal sequence from residue 1 to residue M is cleaved, would have the residues from residue M+1 to residue N remaining. Further as used herein, a "mature" form of a polypeptide or protein may arise from a step of post-translational modification other than a proteolytic cleavage event. Such additional processes include, by way of non-limiting example, glycosylation, myristylation, or phosphorylation. In general, a

mature polypeptide or protein may result from the operation of only one of these processes, or a combination of any of them.

A nucleic acid molecule of the present invention, *e.g.*, a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:1, 2, 4, or 5, or a complement of any of these
5 nucleotide sequences, can be isolated using standard molecular biology techniques and the sequence information provided herein. Using all or a portion of the nucleic acid sequences of SEQ ID NO:1, 2, 4, or 5 as a hybridization probe, FCTR_X molecules can be isolated using standard hybridization and cloning techniques (*e.g.*, as described in Sambrook *et al.*, eds., MOLECULAR CLONING: A LABORATORY MANUAL 2nd Ed., Cold Spring Harbor Laboratory
10 Press, Cold Spring Harbor, NY, 1989; and Ausubel, *et al.*, eds., CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, New York, NY, 1993.)

A nucleic acid of the invention can be amplified using cDNA, mRNA or alternatively, genomic DNA, as a template and appropriate oligonucleotide primers according to standard PCR amplification techniques. The nucleic acid so amplified can be cloned into an
15 appropriate vector and characterized by DNA sequence analysis. Furthermore, oligonucleotides corresponding to FCTR_X nucleotide sequences can be prepared by standard synthetic techniques, *e.g.*, using an automated DNA synthesizer.

In another embodiment, an isolated nucleic acid molecule of the invention comprises a nucleic acid molecule that is a complement of the nucleotide sequence shown in SEQ ID
20 NO:1, 2, 4, or 5. In another embodiment, an isolated nucleic acid molecule of the invention comprises a nucleic acid molecule that is a complement of the nucleotide sequence shown in SEQ ID NO:1, 2, 4, or 5, or a portion of this nucleotide sequence. A nucleic acid molecule that is complementary to the nucleotide sequence shown in SEQ ID NO:1, 2, 4, or 5 is one that is sufficiently complementary to the nucleotide sequence shown in SEQ ID NO:1, 2, 4, or
25 5 that it can hydrogen bond with little or no mismatches to the nucleotide sequence shown in SEQ ID NO:1, 2, 4, or 5 thereby forming a stable duplex..

Moreover, the nucleic acid molecule of the invention can comprise only a portion of the nucleic acid sequence of SEQ ID NO:1, 2, 4, or 5, *e.g.*, a fragment that can be used as a probe or primer or a fragment encoding a biologically active portion of an FCTR_X.
30 Fragments provided herein are defined as sequences of at least 6 (contiguous) nucleic acids or at least 4 (contiguous) amino acids, a length sufficient to allow for specific hybridization in the case of nucleic acids or for specific recognition of an epitope in the case of amino acids, respectively, and are at most some portion less than a full length sequence. Fragments may be derived from any contiguous portion of a nucleic acid or amino acid sequence of choice.

Derivatives are nucleic acid sequences or amino acid sequences formed from the native compounds either directly or by modification or partial substitution. Analogs are nucleic acid sequences or amino acid sequences that have a structure similar to, but not identical to, the native compound but differs from it in respect to certain components or side chains. Analogs
5 may be synthetic or from a different evolutionary origin and may have a similar or opposite metabolic activity compared to wild type. Homologs are nucleic acid sequences or amino acid sequences of a particular gene that are derived from different species.

Derivatives and analogs may be full length or other than full length, if the derivative or analog contains a modified nucleic acid or amino acid, as described below. Derivatives or
10 analogs of the nucleic acids or proteins of the invention include, but are not limited to, molecules comprising regions that are substantially homologous to the nucleic acids or proteins of the invention, in various embodiments, by at least about 30%, 50%, 70%, 80%, or 95% identity (with a preferred identity of 80-95%) over a nucleic acid or amino acid sequence of identical size or when compared to an aligned sequence in which the alignment is done by a
15 computer homology program known in the art, or whose encoding nucleic acid is capable of hybridizing to the complement of a sequence encoding the aforementioned proteins under stringent, moderately stringent, or low stringent conditions. *See e.g.* Ausubel, *et al.*, CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, New York, NY, 1993, and below.

The nucleotide sequence determined from the cloning of a human FCTR_X gene allows
20 for the generation of probes and primers designed for use in identifying and/or cloning FCTR_X homologues in other cell types, *e.g.* from other tissues, as well as FCTR_X homologues from other mammals. The probe/primer typically comprises substantially purified oligonucleotide. The oligonucleotide typically comprises a region of nucleotide sequence that hybridizes under stringent conditions to at least about 12, 25, 50, 100, 150, 200,
25 250, 300, 350 or 400 consecutive sense strand nucleotide sequence of SEQ ID NO:1, 2, 4, or 5 or an anti-sense strand nucleotide sequence of SEQ ID NO:1, 2, 4, or 5 or of a naturally occurring mutant of one or more of these sequences.

Probes based on the human FCTR_X nucleotide sequence can be used to detect transcripts or genomic sequences encoding the same or homologous proteins. In various
30 embodiments, the probe further comprises a label group attached thereto, *e.g.* the label group can be a radioisotope, a fluorescent compound, an enzyme, or an enzyme co-factor. Such probes can be used as a part of a diagnostic test kit for identifying cells or tissue which misexpress an FCTR_X protein, such as by measuring a level of an FCTR_X-encoding nucleic

acid in a sample of cells from a subject *e.g.*, detecting FCTR_X mRNA levels or determining whether a genomic FCTR_X gene has been mutated or deleted.

5 "A polypeptide having a biologically active portion of FCTR_X" refers to polypeptides exhibiting activity similar, but not necessarily identical to, an activity of a polypeptide of the present invention, including mature forms, as measured in a particular biological assay, with or without dose dependency. A nucleic acid fragment encoding a "biologically active portion of FCTR_X" can be prepared by isolating a portion of SEQ ID NO:1, 2, 4, or 5 that encodes a polypeptide having a FCTR_X biological activity (the biological activities of the FCTR_X proteins are described above), expressing the encoded portion of FCTR_X protein (*e.g.*, by
10 recombinant expression *in vitro*) and assessing the activity of the encoded portion of FCTR_X.

FCTR_X variants

The invention further encompasses nucleic acid molecules that differ from the nucleotide sequence shown in SEQ ID NO:1, 2, 4, or 5 due to degeneracy of the genetic code
15 and thus encode the same FCTR_X protein as that encoded by the nucleotide sequence shown in SEQ ID NO:1, 2, 4, or 5. In another embodiment, an isolated nucleic acid molecule of the invention has a nucleotide sequence encoding a protein having an amino acid sequence shown in SEQ ID NO: 3 or 6.

In addition to the human FCTR_X nucleotide sequence shown in SEQ ID NO:1, 2, 4, or
20 5 it will be appreciated by those skilled in the art that DNA sequence polymorphisms that lead to changes in the amino acid sequences of FCTR_X may exist within a population (*e.g.*, the human population). Such genetic polymorphism in the FCTR_X gene may exist among individuals within a population due to natural allelic variation. As used herein, the terms "gene" and "recombinant gene" refer to nucleic acid molecules comprising an open reading
25 frame encoding an FCTR_X protein, preferably a mammalian FCTR_X protein. Such natural allelic variations can typically result in 1-5% variance in the nucleotide sequence of the FCTR_X gene. Any and all such nucleotide variations and resulting amino acid polymorphisms in FCTR_X that are the result of natural allelic variation and that do not alter the functional activity of FCTR_X are intended to be within the scope of the invention.

30 Moreover, nucleic acid molecules encoding FCTR_X proteins from other species, and thus that have a nucleotide sequence that differs from the human sequence of SEQ ID NO:1, 2, 4, or 5 are intended to be within the scope of the invention. Nucleic acid molecules corresponding to natural allelic variants and homologues of the FCTR_X cDNAs of the invention can be isolated based on their homology to the human FCTR_X nucleic acids

disclosed herein using the human cDNAs, or a portion thereof, as a hybridization probe according to standard hybridization techniques under stringent hybridization conditions. For example, a soluble human FCTR_X cDNA can be isolated based on its homology to human membrane-bound FCTR_X cDNA. Likewise, a membrane-bound human FCTR_X cDNA can
5 be isolated based on its homology to soluble human FCTR_X cDNA.

Accordingly, in another embodiment, an isolated nucleic acid molecule of the invention is at least 6 nucleotides in length and hybridizes under stringent conditions to the nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:1, 2, 4, or 5. In another embodiment, the nucleic acid is at least 10, 25, 50, 100, 250 or 500 nucleotides in
10 length. In another embodiment, an isolated nucleic acid molecule of the invention hybridizes to the coding region. As used herein, the term "hybridizes under stringent conditions" is intended to describe conditions for hybridization and washing under which nucleotide sequences at least 60% homologous to each other typically remain hybridized to each other.

Homologs (*i.e.*, nucleic acids encoding FCTR_X proteins derived from species other
15 than human) or other related sequences (*e.g.*, paralogs) can be obtained by low, moderate or high stringency hybridization with all or a portion of the particular human sequence as a probe using methods well known in the art for nucleic acid hybridization and cloning.

Stringent conditions are known to those skilled in the art and can be found in CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6.
20 Preferably, the conditions are such that sequences at least about 65%, 70%, 75%, 85%, 90%, 95%, 98%, or 99% homologous to each other typically remain hybridized to each other. A non-limiting example of stringent hybridization conditions are hybridization in a high salt buffer comprising 6X SSC, 50 mM Tris-HCl (pH 7.5), 1 mM EDTA, 0.02% PVP, 0.02% Ficoll, 0.02% BSA, and 500 mg/ml denatured salmon sperm DNA at 65°C, followed by one or
25 more washes in 0.2X SSC, 0.01% BSA at 50°C. An isolated nucleic acid molecule of the invention that hybridizes under stringent conditions to the sequence of SEQ ID NO:1, 2, 4, or 5 corresponds to a naturally-occurring nucleic acid molecule. As used herein, a "naturally-occurring" nucleic acid molecule refers to an RNA or DNA molecule having a nucleotide sequence that occurs in nature (*e.g.*, encodes a natural protein).

30 In a second embodiment, a nucleic acid sequence that is hybridizable to the nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:1, 2, 4, or 5, or fragments, analogs or derivatives thereof, under conditions of moderate stringency is provided. A non-limiting example of moderate stringency hybridization conditions are hybridization in 6X SSC, 5X Denhardt's solution, 0.5% SDS and 100 mg/ml denatured salmon sperm DNA at

55°C, followed by one or more washes in 1X SSC, 0.1% SDS at 37°C. Other conditions of moderate stringency that may be used are well-known in the art. See, *e.g.*, Ausubel *et al.* (eds.), 1993, CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, NY, and Kriegler, 1990, GENE TRANSFER AND EXPRESSION, A LABORATORY MANUAL, Stockton Press, NY.

In a third embodiment, a nucleic acid that is hybridizable to the nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:1, 2, 4, or 5 or fragments, analogs or derivatives thereof, under conditions of low stringency, is provided. A non-limiting example of low stringency hybridization conditions are hybridization in 35% formamide, 5X SSC, 50 mM Tris-HCl (pH 7.5), 5 mM EDTA, 0.02% PVP, 0.02% Ficoll, 0.2% BSA, 100 mg/ml denatured salmon sperm DNA, 10% (wt/vol) dextran sulfate at 40°C, followed by one or more washes in 2X SSC, 25 mM Tris-HCl (pH 7.4), 5 mM EDTA, and 0.1% SDS at 50°C. Other conditions of low stringency that may be used are well known in the art (*e.g.*, as employed for cross-species hybridizations). See, *e.g.*, Ausubel *et al.* (eds.), 1993, CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, NY, and Kriegler, 1990, GENE TRANSFER AND EXPRESSION, A LABORATORY MANUAL, Stockton Press, NY; Shilo and Weinberg, 1981, *Proc Natl Acad Sci USA* 78: 6789-6792.

Conservative mutations

In addition to naturally-occurring allelic variants of the FCTR_X sequence that may exist in the population, the skilled artisan will further appreciate that changes can be introduced by mutation into the nucleotide sequence of SEQ ID NO:1, 2, 4, or 5 thereby leading to changes in the amino acid sequence of the encoded FCTR_X protein, without altering the functional ability of the FCTR_X protein. For example, nucleotide substitutions leading to amino acid substitutions at "non-essential" amino acid residues can be made in the sequence of SEQ ID NO:1, 2, 4, or 5. A "non-essential" amino acid residue is a residue that can be altered from the wild-type sequence of FCTR_X without altering the biological activity, whereas an "essential" amino acid residue is required for biological activity. For example, amino acid residues that are conserved among the FCTR_X proteins of the present invention, are predicted to be particularly unamenable to alteration.

Another aspect of the invention pertains to nucleic acid molecules encoding FCTR_X proteins that contain changes in amino acid residues that are not essential for activity. Such FCTR_X proteins differ in amino acid sequence from SEQ ID NO:3 or 6, yet retain biological activity. In one embodiment, the isolated nucleic acid molecule comprises a nucleotide

sequence encoding a protein, wherein the protein comprises an amino acid sequence at least about 45% homologous to the amino acid sequence of SEQ ID NO:3 or 6. Preferably, the protein encoded by the nucleic acid molecule is at least about 60% homologous to SEQ ID NO:3 or 6, more preferably at least about 70% homologous to SEQ ID NO:3 or 6, still more preferably at least about 80% homologous to SEQ ID NO:3 or 6, even more preferably at least about 90% homologous to SEQ ID NO:3 or 6, and most preferably at least about 95% homologous to SEQ ID NO:3 or 6.

An isolated nucleic acid molecule encoding a FCTR_X protein homologous to the protein of SEQ ID NO:3 or 6 can be created by introducing one or more nucleotide substitutions, additions or deletions into the nucleotide sequence of SEQ ID NO:1, 2, 4, or 5 such that one or more amino acid substitutions, additions or deletions are introduced into the encoded protein.

Mutations can be introduced into SEQ ID NO:1, 2, 4, or 5 by standard techniques, such as site-directed mutagenesis and PCR-mediated mutagenesis. Preferably, conservative amino acid substitutions are made at one or more predicted non-essential amino acid residues. A "conservative amino acid substitution" is one in which the amino acid residue is replaced with an amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have been defined in the art. These families include amino acids with basic side chains (*e.g.*, lysine, arginine, histidine), acidic side chains (*e.g.*, aspartic acid, glutamic acid), uncharged polar side chains (*e.g.*, glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine), nonpolar side chains (*e.g.*, alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (*e.g.*, threonine, valine, isoleucine) and aromatic side chains (*e.g.*, tyrosine, phenylalanine, tryptophan, histidine). Thus, a predicted nonessential amino acid residue in FCTR_X is replaced with another amino acid residue from the same side chain family. Alternatively, in another embodiment, mutations can be introduced randomly along all or part of a FCTR_X coding sequence, such as by saturation mutagenesis, and the resultant mutants can be screened for FCTR_X biological activity to identify mutants that retain activity. Following mutagenesis of SEQ ID NO:1, 2, 4, or 5, the encoded protein can be expressed by any recombinant technology known in the art and the activity of the protein can be determined.

In one embodiment, a mutant FCTR_X protein can be assayed for (1) the ability to form protein:protein interactions with other FCTR_X proteins, other cell-surface proteins, or biologically active portions thereof, (2) complex formation between a mutant FCTR_X protein

and a FCTR_X ligand; (3) the ability of a mutant FCTR_X protein to bind to an intracellular target protein or biologically active portion thereof; (*e.g.* avidin proteins).

Antisense

5 Another aspect of the invention pertains to isolated antisense nucleic acid molecules that are hybridizable to or complementary to the nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:1, 2, 4, or 5 or fragments, analogs or derivatives thereof. An "antisense" nucleic acid comprises a nucleotide sequence that is complementary to a "sense" nucleic acid encoding a protein, *e.g.*, complementary to the coding strand of a
10 double-stranded cDNA molecule or complementary to an mRNA sequence. In specific aspects, antisense nucleic acid molecules are provided that comprise a sequence complementary to at least about 10, 25, 50, 100, 250 or 500 nucleotides or an entire FCTR_X coding strand, or to only a portion thereof. Nucleic acid molecules encoding fragments, homologs, derivatives and analogs of an FCTR_X protein of SEQ ID NO:3 or 6, or antisense
15 nucleic acids complementary to an FCTR_X nucleic acid sequence of SEQ ID NO:1, 2, 4, or 5 are additionally provided.

 In one embodiment, an antisense nucleic acid molecule is antisense to a "coding region" of the coding strand of a nucleotide sequence encoding FCTR_X. The term "coding region" refers to the region of the nucleotide sequence comprising codons which are translated
20 into amino acid residues. In another embodiment, the antisense nucleic acid molecule is antisense to a "noncoding region" of the coding strand of a nucleotide sequence encoding FCTR_X. The term "noncoding region" refers to 5' and 3' sequences which flank the coding region that are not translated into amino acids (*i.e.*, also referred to as 5' and 3' untranslated regions).

25 Given the coding strand sequences encoding FCTR_X disclosed herein (*e.g.*, SEQ ID NO:1, 2, 4, or 5), antisense nucleic acids of the invention can be designed according to the rules of Watson and Crick or Hoogsteen base pairing. The antisense nucleic acid molecule can be complementary to the entire coding region of FCTR_X mRNA, but more preferably is an oligonucleotide that is antisense to only a portion of the coding or noncoding region of
30 FCTR_X mRNA. For example, the antisense oligonucleotide can be complementary to the region surrounding the translation start site of FCTR_X mRNA. An antisense oligonucleotide can be, for example, about 5, 10, 15, 20, 25, 30, 35, 40, 45 or 50 nucleotides in length. An antisense nucleic acid of the invention can be constructed using chemical synthesis or enzymatic ligation reactions using procedures known in the art. For example, an antisense

nucleic acid (e.g., an antisense oligonucleotide) can be chemically synthesized using naturally occurring nucleotides or variously modified nucleotides designed to increase the biological stability of the molecules or to increase the physical stability of the duplex formed between the antisense and sense nucleic acids, e.g., phosphorothioate derivatives and acridine substituted nucleotides can be used.

Examples of modified nucleotides that can be used to generate the antisense nucleic acid include: 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine. Alternatively, the antisense nucleic acid can be produced biologically using an expression vector into which a nucleic acid has been subcloned in an antisense orientation (i.e., RNA transcribed from the inserted nucleic acid will be of an antisense orientation to a target nucleic acid of interest, described further in the following subsection).

The antisense nucleic acid molecules of the invention are typically administered to a subject or generated *in situ* such that they hybridize with or bind to cellular mRNA and/or genomic DNA encoding a FCTR protein to thereby inhibit expression of the protein, e.g., by inhibiting transcription and/or translation. The hybridization can be by conventional nucleotide complementarity to form a stable duplex, or, for example, in the case of an antisense nucleic acid molecule that binds to DNA duplexes, through specific interactions in the major groove of the double helix. An example of a route of administration of antisense nucleic acid molecules of the invention includes direct injection at a tissue site. Alternatively, antisense nucleic acid molecules can be modified to target selected cells and then administered systemically. For example, for systemic administration, antisense molecules can be modified such that they specifically bind to receptors or antigens expressed on a selected cell surface, e.g., by linking the antisense nucleic acid molecules to peptides or antibodies that bind to cell surface receptors or antigens. The antisense nucleic acid molecules can also be delivered to

cells using the vectors described herein. To achieve sufficient intracellular concentrations of antisense molecules, vector constructs in which the antisense nucleic acid molecule is placed under the control of a strong pol II or pol III promoter are preferred.

In yet another embodiment, the antisense nucleic acid molecule of the invention is an α -anomeric nucleic acid molecule. An α -anomeric nucleic acid molecule forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual β -units, the strands run parallel to each other (Gaultier *et al.* (1987) *Nucleic Acids Res* 15: 6625-6641). The antisense nucleic acid molecule can also comprise a 2'-o-methylribonucleotide (Inoue *et al.* (1987) *Nucleic Acids Res* 15: 6131-6148) or a chimeric RNA-DNA analogue (Inoue *et al.* (1987) *FEBS Lett* 215: 327-330).

Ribozymes and PNA moieties

In still another embodiment, an antisense nucleic acid of the invention is a ribozyme. Ribozymes are catalytic RNA molecules with ribonuclease activity that are capable of cleaving a single-stranded nucleic acid, such as an mRNA, to which they have a complementary region. Thus, ribozymes (*e.g.*, hammerhead ribozymes (described in Haselhoff and Gerlach (1988) *Nature* 334:585-591)) can be used to catalytically cleave FCTR_X mRNA transcripts to thereby inhibit translation of FCTR_X mRNA. A ribozyme having specificity for an FCTR_X-encoding nucleic acid can be designed based upon the nucleotide sequence of an FCTR_X cDNA disclosed herein (*i.e.*, SEQ ID NO:1, 2, 4, or 5). For example, a derivative of a Tetrahymena L-19 IVS RNA can be constructed in which the nucleotide sequence of the active site is complementary to the nucleotide sequence to be cleaved in a FCTR_X-encoding mRNA. See, *e.g.*, Cech *et al.* U.S. Pat. No. 4,987,071; and Cech *et al.* U.S. Pat. No. 5,116,742. Alternatively, FCTR_X mRNA can be used to select a catalytic RNA having a specific ribonuclease activity from a pool of RNA molecules. See, *e.g.*, Bartel *et al.*, (1993) *Science* 261:1411-1418.

Alternatively, FCTR_X gene expression can be inhibited by targeting nucleotide sequences complementary to the regulatory region of the FCTR_X (*e.g.*, the FCTR_X promoter and/or enhancers) to form triple helical structures that prevent transcription of the FCTR_X gene in target cells. See generally, Helene. (1991) *Anticancer Drug Des.* 6: 569-84; Helene. *et al.* (1992) *Ann. N.Y. Acad. Sci.* 660:27-36; and Maher (1992) *Bioassays* 14: 807-15.

In various embodiments, the nucleic acids of FCTR_X can be modified at the base moiety, sugar moiety or phosphate backbone to improve, *e.g.*, the stability, hybridization, or solubility of the molecule. For example, the deoxyribose phosphate backbone of the nucleic

acids can be modified to generate peptide nucleic acids (see Hyrup *et al.* (1996) *Bioorg Med Chem* 4: 5-23). As used herein, the terms "peptide nucleic acids" or "PNAs" refer to nucleic acid mimics, *e.g.*, DNA mimics, in which the deoxyribose phosphate backbone is replaced by a pseudopeptide backbone and only the four natural nucleobases are retained. The neutral backbone of PNAs has been shown to allow for specific hybridization to DNA and RNA under conditions of low ionic strength. The synthesis of PNA oligomers can be performed using standard solid phase peptide synthesis protocols as described in Hyrup *et al.* (1996) above; Perry-O'Keefe *et al.* (1996) *PNAS* 93: 14670-675.

PNAs of FCTR_X can be used in therapeutic and diagnostic applications. For example, PNAs can be used as antisense or antigene agents for sequence-specific modulation of gene expression by, *e.g.*, inducing transcription or translation arrest or inhibiting replication. PNAs of FCTR_X can also be used, *e.g.*, in the analysis of single base pair mutations in a gene by, *e.g.*, PNA directed PCR clamping; as artificial restriction enzymes when used in combination with other enzymes, *e.g.*, S1 nucleases (Hyrup B. (1996) above); or as probes or primers for DNA sequence and hybridization (Hyrup *et al.* (1996), above; Perry-O'Keefe (1996), above).

In another embodiment, PNAs of FCTR_X can be modified, *e.g.*, to enhance their stability or cellular uptake, by attaching lipophilic or other helper groups to PNA, by the formation of PNA-DNA chimeras, or by the use of liposomes or other techniques of drug delivery known in the art. For example, PNA-DNA chimeras of FCTR_X can be generated that may combine the advantageous properties of PNA and DNA. Such chimeras allow DNA recognition enzymes, *e.g.*, RNase H and DNA polymerases, to interact with the DNA portion while the PNA portion would provide high binding affinity and specificity. PNA-DNA chimeras can be linked using linkers of appropriate lengths selected in terms of base stacking, number of bonds between the nucleobases, and orientation (Hyrup (1996) above). The synthesis of PNA-DNA chimeras can be performed as described in Hyrup (1996) above and Finn *et al.* (1996) *Nucl Acids Res* 24: 3357-63. For example, a DNA chain can be synthesized on a solid support using standard phosphoramidite coupling chemistry, and modified nucleoside analogs, *e.g.*, 5'-(4-methoxytrityl)amino-5'-deoxy-thymidine phosphoramidite, can be used between the PNA and the 5' end of DNA (Mag *et al.* (1989) *Nucl Acid Res* 17: 5973-88). PNA monomers are then coupled in a stepwise manner to produce a chimeric molecule with a 5' PNA segment and a 3' DNA segment (Finn *et al.* (1996) above). Alternatively, chimeric molecules can be synthesized with a 5' DNA segment and a 3' PNA segment. See, Petersen *et al.* (1975) *Bioorg Med Chem Lett* 5: 1119-11124.

In other embodiments, the oligonucleotide may include other appended groups such as peptides (e.g., for targeting host cell receptors *in vivo*), or agents facilitating transport across the cell membrane (see, e.g., Letsinger *et al.*, 1989, *Proc. Natl. Acad. Sci. U.S.A.* 86:6553-6556; Lemaitre *et al.*, 1987, *Proc. Natl. Acad. Sci.* 84:648-652; PCT Publication No. W088/09810) or the blood-brain barrier (see, e.g., PCT Publication No. W089/10134). In addition, oligonucleotides can be modified with hybridization triggered cleavage agents (See, e.g., Krol *et al.*, 1988, *BioTechniques* 6:958-976) or intercalating agents. (See, e.g., Zon, 1988, *Pharm. Res.* 5: 539-549). To this end, the oligonucleotide may be conjugated to another molecule, e.g., a peptide, a hybridization triggered cross-linking agent, a transport agent, a hybridization-triggered cleavage agent, etc.

FCTR proteins

One aspect of the invention pertains to isolated FCTR proteins, and biologically active portions thereof, or derivatives, fragments, analogs or homologs thereof. Also provided are polypeptide fragments suitable for use as immunogens to raise anti-FCTR antibodies. In one embodiment, native FCTR proteins can be isolated from cells or tissue sources by an appropriate purification scheme using standard protein purification techniques. In another embodiment, FCTR proteins are produced by recombinant DNA techniques. Alternative to recombinant expression, a FCTR protein or polypeptide can be synthesized chemically using standard peptide synthesis techniques.

An "isolated" or "purified" protein or biologically active portion thereof is substantially free of cellular material or other contaminating proteins from the cell or tissue source from which the FCTR protein is derived, or substantially free from chemical precursors or other chemicals when chemically synthesized. The language "substantially free of cellular material" includes preparations of FCTR protein in which the protein is separated from cellular components of the cells from which it is isolated or recombinantly produced. In one embodiment, the language "substantially free of cellular material" includes preparations of FCTR protein having less than about 30% (by dry weight) of non-FCTR protein (also referred to herein as a "contaminating protein"), more preferably less than about 20% of non-FCTR protein, still more preferably less than about 10% of non-FCTR protein, and most preferably less than about 5% non-FCTR protein. When the FCTR protein or biologically active portion thereof is recombinantly produced, it is also preferably substantially free of culture medium, *i.e.*, culture medium represents less than about 20%,

more preferably less than about 10%, and most preferably less than about 5% of the volume of the protein preparation.

The language "substantially free of chemical precursors or other chemicals" includes preparations of FCTR_X protein in which the protein is separated from chemical precursors or other chemicals that are involved in the synthesis of the protein. In one embodiment, the language "substantially free of chemical precursors or other chemicals" includes preparations of FCTR_X protein having less than about 30% (by dry weight) of chemical precursors or non-FCTR_X chemicals, more preferably less than about 20% chemical precursors or non-FCTR_X chemicals, still more preferably less than about 10% chemical precursors or non-FCTR_X chemicals, and most preferably less than about 5% chemical precursors or non-FCTR_X chemicals.

Biologically active portions of an FCTR_X protein include peptides comprising amino acid sequences sufficiently homologous to or derived from the amino acid sequence of the FCTR_X protein, *e.g.*, the amino acid sequence shown in SEQ ID NO:3 or 6, that include fewer amino acids than the full length FCTR_X proteins, and exhibit at least one activity of a FCTR_X protein. Typically, biologically active portions comprise a domain or motif with at least one activity of the FCTR_X protein. A biologically active portion of an FCTR_X protein can be a polypeptide which is, for example, 10, 25, 50, 100 or more amino acids in length.

Moreover, other biologically active portions, in which other regions of the protein are deleted, can be peppered by recombinant techniques and evaluated for one or more of the functional activities of a native FCTR_X protein.

In an embodiment, the FCTR_X protein has an amino acid sequence shown in SEQ ID NO:3 or 6. In other embodiments, the FCTR_X protein is substantially homologous to SEQ ID NO:3 or 6, and retains the functional activity of the protein of SEQ ID NO:3 or 6, yet differs in amino acid sequence due to natural allelic variation or mutagenesis, as described in detail below. Accordingly, in another embodiment, the FCTR_X protein is a protein that comprises an amino acid sequence at least about 45% homologous to the amino acid sequence of SEQ ID NO:3 or 6 and retains the functional activity of the FCTR_X proteins of SEQ ID NO:3 or 6.

Determining homology between two or more sequences

To determine the percent homology of two amino acid sequences or of two nucleic acids, the sequences are aligned for optimal comparison purposes (*e.g.*, gaps can be introduced in the sequence of a first amino acid or nucleic acid sequence for optimal alignment with a second amino or nucleic acid sequence). The amino acid residues or nucleotides at

corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are homologous at that position (*i.e.*, as used herein amino acid or nucleic acid "homology" is equivalent to amino acid or nucleic acid "identity").

The nucleic acid sequence homology may be determined as the degree of identity between two sequences. The homology may be determined using computer programs known in the art, such as GAP software provided in the GCG program package. See, *Needleman and Wunsch* 1970 *J Mol Biol* 48: 443-453. Using GCG GAP software with the following settings for nucleic acid sequence comparison: GAP creation penalty of 5.0 and GAP extension penalty of 0.3, the coding region of the analogous nucleic acid sequences referred to above exhibits a degree of identity preferably of at least 70%, 75%, 80%, 85%, 90%, 95%, 98%, or 99%, with the CDS (encoding) part of the DNA sequence shown in SEQ ID NO:1, 2, 4, or 5.

The term "sequence identity" refers to the degree to which two polynucleotide or polypeptide sequences are identical on a residue-by-residue basis over a particular region of comparison. The term "percentage of sequence identity" is calculated by comparing two optimally aligned sequences over that region of comparison, determining the number of positions at which the identical nucleic acid base (*e.g.*, A, T, C, G, U, or I, in the case of nucleic acids) occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the region of comparison (*i.e.*, the window size), and multiplying the result by 100 to yield the percentage of sequence identity. The term "substantial identity" as used herein denotes a characteristic of a polynucleotide sequence, wherein the polynucleotide comprises a sequence that has at least 80 percent sequence identity, preferably at least 85 percent identity and often 90 to 95 percent sequence identity, more usually at least 99 percent sequence identity as compared to a reference sequence over a comparison region.

Chimeric and fusion proteins

The invention also provides FCTR_X chimeric or fusion proteins. As used herein, an FCTR_X "chimeric protein" or "fusion protein" comprises an FCTR_X polypeptide operatively linked to a non-FCTR_X polypeptide. An "FCTR_X polypeptide" refers to a polypeptide having an amino acid sequence corresponding to FCTR_X, whereas a "non-FCTR_X polypeptide" refers to a polypeptide having an amino acid sequence corresponding to a protein that is not substantially homologous to the FCTR_X protein, *e.g.*, a protein that is different from the

FCTR_X protein and that is derived from the same or a different organism. Within an FCTR_X fusion protein the FCTR_X polypeptide can correspond to all or a portion of an FCTR_X protein. In one embodiment, an FCTR_X fusion protein comprises at least one biologically active portion of an FCTR_X protein. In another embodiment, an FCTR_X fusion protein
5 comprises at least two biologically active portions of an FCTR_X protein. In yet another embodiment, an FCTR_X fusion protein comprises at least three biologically active portions of an FCTR_X protein. Within the fusion protein, the term "operatively linked" is intended to indicate that the FCTR_X polypeptide and the non-FCTR_X polypeptide are fused in-frame to each other. The non-FCTR_X polypeptide can be fused to the N-terminus or C-terminus of the
10 FCTR_X polypeptide.

In yet another embodiment, the fusion protein is a GST-FCTR_X fusion protein in which the FCTR_X sequences are fused to the C-terminus of the GST (*i.e.*, glutathione S-transferase) sequences. Such fusion proteins can facilitate the purification of recombinant FCTR_X.

15 In another embodiment, the fusion protein is an FCTR_X protein containing a heterologous signal sequence at its N-terminus. For example, the native FCTR_X signal sequence can be removed and replaced with a signal sequence from another protein. In certain host cells (*e.g.*, mammalian host cells), expression and/or secretion of FCTR_X can be increased through use of a heterologous signal sequence.

20 In yet another embodiment, the fusion protein is an FCTR_X-immunoglobulin fusion protein in which the FCTR_X are fused to sequences derived from a member of the immunoglobulin protein family. The FCTR_X-immunoglobulin fusion proteins of the invention can be incorporated into pharmaceutical compositions and administered to a subject to inhibit an interaction between an FCTR_X ligand and an FCTR_X protein on the surface of a
25 cell, to thereby suppress FCTR_X-mediated signal transduction *in vivo*. The FCTR_X-immunoglobulin fusion proteins can be used to affect the bioavailability of an FCTR_X cognate ligand. Inhibition of the FCTR_X ligand/FCTR_X interaction are useful therapeutically for both the treatment of proliferative and differentiative disorders, as well as modulating (*e.g.* promoting or inhibiting) cell survival. Moreover, the
30 FCTR_X-immunoglobulin fusion proteins of the invention can be used as immunogens to produce anti-FCTR_X antibodies in a subject, to purify FCTR_X ligands, and in screening assays to identify molecules that inhibit the interaction of FCTR_X with an FCTR_X ligand.

An FCTR_X chimeric or fusion protein of the invention can be produced by standard recombinant DNA techniques. For example, DNA fragments coding for the different

polypeptide sequences are ligated together in-frame in accordance with conventional techniques, *e.g.*, by employing blunt-ended or stagger-ended termini for ligation, restriction enzyme digestion to provide for appropriate termini, filling-in of cohesive ends as appropriate, alkaline phosphatase treatment to avoid undesirable joining, and enzymatic ligation. In another embodiment, the fusion gene can be synthesized by conventional techniques including automated DNA synthesizers. Alternatively, PCR amplification of gene fragments can be carried out using anchor primers that give rise to complementary overhangs between two consecutive gene fragments that can subsequently be annealed and reamplified to generate a chimeric gene sequence (see, for example, Ausubel *et al.* (eds.) CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, 1992). Moreover, many expression vectors are commercially available that already encode a fusion moiety (*e.g.*, a GST polypeptide). An FCTR_X-encoding nucleic acid can be cloned into such an expression vector such that the fusion moiety is linked in-frame to the FCTR_X protein.

15 FCTR_X agonists and antagonists

The present invention also pertains to variants of the FCTR_X proteins that function as either FCTR_X agonists (mimetics) or as FCTR_X antagonists. Variants of the FCTR_X protein can be generated by mutagenesis, *e.g.*, discrete point mutation or truncation of the FCTR_X protein. An agonist of the FCTR_X protein can retain substantially the same, or a subset of, the biological activities of the naturally occurring form of the FCTR_X protein. An antagonist of the FCTR_X protein can inhibit one or more of the activities of the naturally occurring form of the FCTR_X protein by, for example, competitively binding to a downstream or upstream member of a cellular signaling cascade which includes the FCTR_X protein. Thus, specific biological effects can be elicited by treatment with a variant of limited function. In one embodiment, treatment of a subject with a variant having a subset of the biological activities of the naturally occurring form of the protein has fewer side effects in a subject relative to treatment with the naturally occurring form of the FCTR_X proteins.

Variants of the FCTR_X protein that function as either FCTR_X agonists (mimetics) or as FCTR_X antagonists can be identified by screening combinatorial libraries of mutants, *e.g.*, truncation mutants, of the FCTR_X protein for FCTR_X protein agonist or antagonist activity. In one embodiment, a variegated library of FCTR_X variants is generated by combinatorial mutagenesis at the nucleic acid level and is encoded by a variegated gene library. A variegated library of FCTR_X variants can be produced by, for example, enzymatically ligating a mixture of synthetic oligonucleotides into gene sequences such that a degenerate set of

potential FCTR_X sequences is expressible as individual polypeptides, or alternatively, as a set of larger fusion proteins (*e.g.*, for phage display) containing the set of FCTR_X sequences therein. There are a variety of methods which can be used to produce libraries of potential FCTR_X variants from a degenerate oligonucleotide sequence. Chemical synthesis of a
5 degenerate gene sequence can be performed in an automatic DNA synthesizer, and the synthetic gene then ligated into an appropriate expression vector. Use of a degenerate set of genes allows for the provision, in one mixture, of all of the sequences encoding the desired set of potential FCTR_X sequences. Methods for synthesizing degenerate oligonucleotides are known in the art (see, *e.g.*, Narang (1983) *Tetrahedron* 39:3; Itakura *et al.* (1984) *Annu Rev*
10 *Biochem* 53:323; Itakura *et al.* (1984) *Science* 198:1056; Ike *et al.* (1983) *Nucl Acid Res* 11:477.

Polypeptide libraries

In addition, libraries of fragments of the FCTR_X protein coding sequence can be used
15 to generate a variegated population of FCTR_X fragments for screening and subsequent selection of variants of an FCTR_X protein. In one embodiment, a library of coding sequence fragments can be generated by treating a double stranded PCR fragment of an FCTR_X coding sequence with a nuclease under conditions wherein nicking occurs only about once per molecule, denaturing the double stranded DNA, renaturing the DNA to form double stranded
20 DNA that can include sense/antisense pairs from different nicked products, removing single stranded portions from reformed duplexes by treatment with S1 nuclease, and ligating the resulting fragment library into an expression vector. By this method, an expression library can be derived which encodes N-terminal and internal fragments of various sizes of the FCTR_X protein.

25 Several techniques are known in the art for screening gene products of combinatorial libraries made by point mutations or truncation, and for screening cDNA libraries for gene products having a selected property. Such techniques are adaptable for rapid screening of the gene libraries generated by the combinatorial mutagenesis of FCTR_X proteins. The most widely used techniques, which are amenable to high throughput analysis, for screening large
30 gene libraries typically include cloning the gene library into replicable expression vectors, transforming appropriate cells with the resulting library of vectors, and expressing the combinatorial genes under conditions in which detection of a desired activity facilitates isolation of the vector encoding the gene whose product was detected. Recursive ensemble mutagenesis (REM), a new technique that enhances the frequency of functional mutants in the

libraries, can be used in combination with the screening assays to identify FCTR_X variants (Arkin and Yourvan (1992) PNAS 89:7811-7815; Delgrave *et al.* (1993) Protein Engineering 6:327-331).

5 Anti-FCTR_X antibodies

An isolated FCTR_X protein, or a portion or fragment thereof, can be used as an immunogen to generate antibodies that bind FCTR_X using standard techniques for polyclonal and monoclonal antibody preparation. The full-length FCTR_X protein can be used or, alternatively, the invention provides antigenic peptide fragments of FCTR_X for use as
10 immunogens. The antigenic peptide of FCTR_X comprises at least 8 amino acid residues of the amino acid sequence shown in SEQ ID NO:3 or 6 and encompasses an epitope of FCTR_X such that an antibody raised against the peptide forms a specific immune complex with FCTR_X. Preferably, the antigenic peptide comprises at least 10 amino acid residues, more preferably at least 15 amino acid residues, even more preferably at least 20 amino acid
15 residues, and most preferably at least 30 amino acid residues. Preferred epitopes encompassed by the antigenic peptide are regions of FCTR_X that are located on the surface of the protein, *e.g.*, hydrophilic regions.

As disclosed herein, FCTR_X protein sequence of SEQ ID NO:3 or 6, or derivatives, fragments, analogs or homologs thereof, may be utilized as immunogens in the generation of
20 antibodies that immunospecifically-bind these protein components. The term "antibody" as used herein refers to immunoglobulin molecules and immunologically active portions of immunoglobulin molecules, *i.e.*, molecules that contain an antigen binding site that specifically binds (immunoreacts with) an antigen, such as FCTR_X. Such antibodies include, but are not limited to, polyclonal, monoclonal, chimeric, single chain, F_{ab} and F_{(ab)²} fragments,
25 and an F_{ab} expression library. In a specific embodiment, antibodies to human FCTR_X proteins are disclosed. Various procedures known within the art may be used for the production of polyclonal or monoclonal antibodies to an FCTR_X protein sequence of SEQ ID NO:3 or 6, or derivative, fragment, analog or homolog thereof.

For the production of polyclonal antibodies, various suitable host animals (*e.g.*, rabbit,
30 goat, mouse or other mammal) may be immunized by injection with the native protein, or a synthetic variant thereof, or a derivative of the foregoing. An appropriate immunogenic preparation can contain, for example, recombinantly expressed FCTR_X protein or a chemically synthesized FCTR_X polypeptide. The preparation can further include an adjuvant. Various adjuvants used to increase the immunological response include, but are not limited to,

Freund's (complete and incomplete), mineral gels (*e.g.*, aluminum hydroxide), surface active substances (*e.g.*, lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, dinitrophenol, etc.), human adjuvants such as *Bacille Calmette-Guerin* and *Corynebacterium parvum*, or similar immunostimulatory agents. If desired, the antibody molecules directed
5 against FCTR_X can be isolated from the mammal (*e.g.*, from the blood) and further purified by well known techniques, such as protein A chromatography to obtain the IgG fraction.

The term "monoclonal antibody" or "monoclonal antibody composition", as used herein, refers to a population of antibody molecules that contain only one species of an antigen binding site capable of immunoreacting with a particular epitope of FCTR_X. A monoclonal
10 antibody composition thus typically displays a single binding affinity for a particular FCTR_X protein with which it immunoreacts. For preparation of monoclonal antibodies directed towards a particular FCTR_X protein, or derivatives, fragments, analogs or homologs thereof, any technique that provides for the production of antibody molecules by continuous cell line culture may be utilized. Such techniques include, but are not limited to, the hybridoma
15 technique (see Kohler & Milstein, 1975 *Nature* 256: 495-497); the trioma technique; the human B-cell hybridoma technique (see Kozbor, *et al.*, 1983 *Immunol Today* 4: 72) and the EBV hybridoma technique to produce human monoclonal antibodies (see Cole, *et al.*, 1985 In: MONOCLONAL ANTIBODIES AND CANCER THERAPY, Alan R. Liss, Inc., pp. 77-96). Human monoclonal antibodies may be utilized in the practice of the present invention and may be
20 produced by using human hybridomas (see Cote, *et al.*, 1983. *Proc Natl Acad Sci USA* 80: 2026-2030) or by transforming human B-cells with Epstein Barr Virus *in vitro* (see Cole, *et al.*, 1985 In: MONOCLONAL ANTIBODIES AND CANCER THERAPY, Alan R. Liss, Inc., pp. 77-96).

According to the invention, techniques can be adapted for the production of single-chain antibodies specific to an FCTR_X protein (see *e.g.*, U.S. Patent No. 4,946,778). In
25 addition, methodologies can be adapted for the construction of F_{ab} expression libraries (see *e.g.*, Huse, *et al.*, 1989 *Science* 246: 1275-1281) to allow rapid and effective identification of monoclonal F_{ab} fragments with the desired specificity for an FCTR_X protein or derivatives, fragments, analogs or homologs thereof. Non-human antibodies can be "humanized" by techniques well known in the art. See *e.g.*, U.S. Patent No. 5,225,539. Antibody fragments
30 that contain the idiotypes to an FCTR_X protein may be produced by techniques known in the art including, but not limited to: (i) an F_{(ab')₂} fragment produced by pepsin digestion of an antibody molecule; (ii) an F_{ab} fragment generated by reducing the disulfide bridges of an F_{(ab')₂} fragment; (iii) an F_{ab} fragment generated by the treatment of the antibody molecule with papain and a reducing agent and (iv) F_v fragments.

Additionally, recombinant anti-FCTR_X antibodies, such as chimeric and humanized monoclonal antibodies, comprising both human and non-human portions, which can be made using standard recombinant DNA techniques, are within the scope of the invention. Such chimeric and humanized monoclonal antibodies can be produced by recombinant DNA

5 techniques known in the art, for example using methods described in PCT International Application No. PCT/US86/02269; European Patent Application No. 184,187; European Patent Application No. 171,496; European Patent Application No. 173,494; PCT International Publication No. WO 86/01533; U.S. Pat. No. 4,816,567; European Patent Application No. 125,023; Better *et al.* (1988) *Science* 240:1041-1043; Liu *et al.* (1987) *PNAS* 84:3439-3443;

10 Liu *et al.* (1987) *J Immunol.* 139:3521-3526; Sun *et al.* (1987) *PNAS* 84:214-218; Nishimura *et al.* (1987) *Cancer Res* 47:999-1005; Wood *et al.* (1985) *Nature* 314:446-449; Shaw *et al.* (1988) *J Natl Cancer Inst* 80:1553-1559; Morrison (1985) *Science* 229:1202-1207; Oi *et al.* (1986) *BioTechniques* 4:214; U.S. Pat. No. 5,225,539; Jones *et al.* (1986) *Nature* 321:552-525; Verhoeyan *et al.* (1988) *Science* 239:1534; and Beidler *et al.* (1988) *J Immunol*

15 141:4053-4060.

In one embodiment, methodologies for the screening of antibodies that possess the desired specificity include, but are not limited to, enzyme-linked immunosorbent assay (ELISA) and other immunologically-mediated techniques known within the art.

Anti-FCTR_X antibodies may be used in methods known within the art relating to the

20 localization and/or quantitation of an FCTR_X protein (*e.g.*, for use in measuring levels of the FCTR_X protein within appropriate physiological samples, for use in diagnostic methods, for use in imaging the protein, and the like). In a given embodiment, antibodies for FCTR_X proteins, or derivatives, fragments, analogs or homologs thereof, that contain the antibody derived binding domain, are utilized as pharmacologically-active compounds [hereinafter

25 "Therapeutics"].

An anti-FCTR_X antibody (*e.g.*, monoclonal antibody) can be used to isolate FCTR_X by standard techniques, such as affinity chromatography or immunoprecipitation. An anti-FCTR_X antibody can facilitate the purification of natural FCTR_X from cells and of recombinantly produced FCTR_X expressed in host cells. Moreover, an anti-FCTR_X antibody

30 can be used to detect FCTR_X protein (*e.g.*, in a cellular lysate or cell supernatant) in order to evaluate the abundance and pattern of expression of the FCTR_X protein. Anti-FCTR_X antibodies can be used diagnostically to monitor protein levels in tissue as part of a clinical testing procedure, *e.g.*, to, for example, determine the efficacy of a given treatment regimen. Detection can be facilitated by coupling (*i.e.*, physically linking) the antibody to a detectable

substance. Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, and radioactive materials. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, β -galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; examples of bioluminescent materials include luciferase, luciferin, and aequorin, and examples of suitable radioactive material include ^{125}I , ^{131}I , ^{35}S or ^3H .

FCTR_X Recombinant Expression Vectors and Host Cells

Another aspect of the invention pertains to vectors, preferably expression vectors, containing a nucleic acid encoding FCTR_X protein, or derivatives, fragments, analogs or homologs thereof. As used herein, the term "vector" refers to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a "plasmid", which refers to a circular double stranded DNA loop into which additional DNA segments can be ligated. Another type of vector is a viral vector, wherein additional DNA segments can be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (*e.g.*, bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (*e.g.*, non-episomal mammalian vectors) are integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to which they are operatively linked. Such vectors are referred to herein as "expression vectors". In general, expression vectors of utility in recombinant DNA techniques are often in the form of plasmids. In the present specification, "plasmid" and "vector" can be used interchangeably as the plasmid is the most commonly used form of vector. However, the invention is intended to include such other forms of expression vectors, such as viral vectors (*e.g.*, replication defective retroviruses, adenoviruses and adeno-associated viruses), which serve equivalent functions.

The recombinant expression vectors of the invention comprise a nucleic acid of the invention in a form suitable for expression of the nucleic acid in a host cell, which means that the recombinant expression vectors include one or more regulatory sequences, selected on the basis of the host cells to be used for expression, that is operatively linked to the nucleic acid

sequence to be expressed. Within a recombinant expression vector, "operably linked" is intended to mean that the nucleotide sequence of interest is linked to the regulatory sequence(s) in a manner that allows for expression of the nucleotide sequence (*e.g.*, in an *in vitro* transcription/translation system or in a host cell when the vector is introduced into the host cell). The term "regulatory sequence" is intended to include promoters, enhancers and other expression control elements (*e.g.*, polyadenylation signals). Such regulatory sequences are described, for example, in Goeddel; GENE EXPRESSION TECHNOLOGY: METHODS IN ENZYMOLOGY 185, Academic Press, San Diego, Calif. (1990). Regulatory sequences include those that direct constitutive expression of a nucleotide sequence in many types of host cell and those that direct expression of the nucleotide sequence only in certain host cells (*e.g.*, tissue-specific regulatory sequences). It will be appreciated by those skilled in the art that the design of the expression vector can depend on such factors as the choice of the host cell to be transformed, the level of expression of protein desired, etc. The expression vectors of the invention can be introduced into host cells to thereby produce proteins or peptides, including fusion proteins or peptides, encoded by nucleic acids as described herein (*e.g.*, FCTR_X proteins, mutant forms of FCTR_X, fusion proteins, etc.).

The recombinant expression vectors of the invention can be designed for expression of FCTR_X in prokaryotic or eukaryotic cells. For example, FCTR_X can be expressed in bacterial cells such as *E. coli*, insect cells (using baculovirus expression vectors) yeast cells or mammalian cells. Suitable host cells are discussed further in Goeddel, GENE EXPRESSION TECHNOLOGY: METHODS IN ENZYMOLOGY 185, Academic Press, San Diego, Calif. (1990). Alternatively, the recombinant expression vector can be transcribed and translated *in vitro*, for example using T7 promoter regulatory sequences and T7 polymerase.

Expression of proteins in prokaryotes is most often carried out in *E. coli* with vectors containing constitutive or inducible promoters directing the expression of either fusion or non-fusion proteins. Fusion vectors add a number of amino acids to a protein encoded therein, usually to the amino terminus of the recombinant protein. Such fusion vectors typically serve three purposes: (1) to increase expression of recombinant protein; (2) to increase the solubility of the recombinant protein; and (3) to aid in the purification of the recombinant protein by acting as a ligand in affinity purification. Often, in fusion expression vectors, a proteolytic cleavage site is introduced at the junction of the fusion moiety and the recombinant protein to enable separation of the recombinant protein from the fusion moiety subsequent to purification of the fusion protein. Such enzymes, and their cognate recognition sequences, include Factor Xa, thrombin and enterokinase. Typical fusion expression vectors include pGEX (Pharmacia

Biotech Inc; Smith and Johnson (1988) *Gene* 67:31-40), pMAL (New England Biolabs, Beverly, Mass.) and pRIT5 (Pharmacia, Piscataway, N.J.) that fuse glutathione S-transferase (GST), maltose E binding protein, or protein A, respectively, to the target recombinant protein.

Examples of suitable inducible non-fusion *E. coli* expression vectors include pTrc
5 (Amrann *et al.*, (1988) *Gene* 69:301-315) and pET 11d (Studier *et al.*, GENE EXPRESSION
TECHNOLOGY: METHODS IN ENZYMOLOGY 185, Academic Press, San Diego, Calif. (1990)
60-89).

One strategy to maximize recombinant protein expression in *E. coli* is to express the
protein in a host bacteria with an impaired capacity to proteolytically cleave the recombinant
10 protein. See, Gottesman, GENE EXPRESSION TECHNOLOGY: METHODS IN ENZYMOLOGY 185,
Academic Press, San Diego, Calif. (1990) 119-128. Another strategy is to alter the nucleic
acid sequence of the nucleic acid to be inserted into an expression vector so that the individual
codons for each amino acid are those preferentially utilized in *E. coli* (Wada *et al.*, (1992)
Nucleic Acids Res. 20:2111-2118). Such alteration of nucleic acid sequences of the invention
15 can be carried out by standard DNA synthesis techniques.

In another embodiment, the FCTR_X expression vector is a yeast expression vector.
Examples of vectors for expression in yeast *S. cerevisiae* include pYepSec1 (Baldari, *et al.*,
(1987) *EMBO J* 6:229-234), pMFa (Kurjan and Herskowitz, (1982) *Cell* 30:933-943), pJRY88
(Schultz *et al.*, (1987) *Gene* 54:113-123), pYES2 (Invitrogen Corporation, San Diego, Calif.),
20 and picZ (Invitrogen Corp, San Diego, Calif.).

Alternatively, FCTR_X can be expressed in insect cells using baculovirus expression
vectors. Baculovirus vectors available for expression of proteins in cultured insect cells (*e.g.*,
SF9 cells) include the pAc series (Smith *et al.* (1983) *Mol Cell Biol* 3:2156-2165) and the pVL
series (Lucklow and Summers (1989) *Virology* 170:31-39).

25 In yet another embodiment, a nucleic acid of the invention is expressed in mammalian
cells using a mammalian expression vector. Examples of mammalian expression vectors
include pCDM8 (Seed (1987) *Nature* 329:840) and pMT2PC (Kaufman *et al.* (1987) *EMBO J*
6: 187-195). When used in mammalian cells, the expression vector's control functions are
often provided by viral regulatory elements. For example, commonly used promoters are
30 derived from polyoma, Adenovirus 2, cytomegalovirus and Simian Virus 40. For other
suitable expression systems for both prokaryotic and eukaryotic cells. See, *e.g.*, Chapters 16
and 17 of Sambrook *et al.*, MOLECULAR CLONING: A LABORATORY MANUAL. 2nd ed., Cold
Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.,
1989.

In another embodiment, the recombinant mammalian expression vector is capable of directing expression of the nucleic acid preferentially in a particular cell type (e.g., tissue-specific regulatory elements are used to express the nucleic acid). Tissue-specific regulatory elements are known in the art. Non-limiting examples of suitable tissue-specific promoters include the albumin promoter (liver-specific; Pinkert *et al.* (1987) *Genes Dev* 1:268-277), lymphoid-specific promoters (Calame and Eaton (1988) *Adv Immunol* 43:235-275), in particular promoters of T cell receptors (Winoto and Baltimore (1989) *EMBO J* 8:729-733) and immunoglobulins (Banerji *et al.* (1983) *Cell* 33:729-740; Queen and Baltimore (1983) *Cell* 33:741-748), neuron-specific promoters (e.g., the neurofilament promoter; Byrne and Ruddle (1989) *PNAS* 86:5473-5477), pancreas-specific promoters (Edlund *et al.* (1985) *Science* 230:912-916), and mammary gland-specific promoters (e.g., milk whey promoter; U.S. Pat. No. 4,873,316 and European Application Publication No. 264,166). Developmentally-regulated promoters are also encompassed, e.g., the murine hox promoters (Kessel and Gruss (1990) *Science* 249:374-379) and the α -fetoprotein promoter (Campes and Tilghman (1989) *Genes Dev* 3:537-546).

The invention further provides a recombinant expression vector comprising a DNA molecule of the invention cloned into the expression vector in an antisense orientation. That is, the DNA molecule is operatively linked to a regulatory sequence in a manner that allows for expression (by transcription of the DNA molecule) of an RNA molecule that is antisense to FCTR_X mRNA. Regulatory sequences operatively linked to a nucleic acid cloned in the antisense orientation can be chosen that direct the continuous expression of the antisense RNA molecule in a variety of cell types, for instance viral promoters and/or enhancers, or regulatory sequences can be chosen that direct constitutive, tissue specific or cell type specific expression of antisense RNA. The antisense expression vector can be in the form of a recombinant plasmid, phagemid or attenuated virus in which antisense nucleic acids are produced under the control of a high efficiency regulatory region, the activity of which can be determined by the cell type into which the vector is introduced. For a discussion of the regulation of gene expression using antisense genes see Weintraub *et al.*, "Antisense RNA as a molecular tool for genetic analysis," Reviews--Trends in Genetics, Vol. 1(1) 1986.

Another aspect of the invention pertains to host cells into which a recombinant expression vector of the invention has been introduced. The terms "host cell" and "recombinant host cell" are used interchangeably herein. It is understood that such terms refer not only to the particular subject cell but to the progeny or potential progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or

environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term as used herein.

5 A host cell can be any prokaryotic or eukaryotic cell. For example, FCTR protein can be expressed in bacterial cells such as *E. coli*, insect cells, yeast or mammalian cells (such as Chinese hamster ovary cells (CHO) or COS cells). Other suitable host cells are known to those skilled in the art.

10 Vector DNA can be introduced into prokaryotic or eukaryotic cells via conventional transformation or transfection techniques. As used herein, the terms "transformation" and "transfection" are intended to refer to a variety of art-recognized techniques for introducing foreign nucleic acid (*e.g.*, DNA) into a host cell, including calcium phosphate or calcium chloride co-precipitation, DEAE-dextran-mediated transfection, lipofection, or electroporation. Suitable methods for transforming or transfecting host cells can be found in Sambrook, *et al.* (MOLECULAR CLONING: A LABORATORY MANUAL. 2nd ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989), and other laboratory manuals.

20 For stable transfection of mammalian cells, it is known that, depending upon the expression vector and transfection technique used, only a small fraction of cells may integrate the foreign DNA into their genome. In order to identify and select these integrants, a gene that encodes a selectable marker (*e.g.*, resistance to antibiotics) is generally introduced into the host cells along with the gene of interest. Various selectable markers include those that confer resistance to drugs, such as G418, hygromycin and methotrexate. Nucleic acid encoding a selectable marker can be introduced into a host cell on the same vector as that encoding FCTR or can be introduced on a separate vector. Cells stably transfected with the introduced nucleic acid can be identified by drug selection (*e.g.*, cells that have incorporated the selectable marker gene will survive, while the other cells die).

25 A host cell of the invention, such as a prokaryotic or eukaryotic host cell in culture, can be used to produce (*i.e.*, express) FCTR protein. Accordingly, the invention further provides methods for producing FCTR protein using the host cells of the invention. In one embodiment, the method comprises culturing the host cell of invention (into which a recombinant expression vector encoding FCTR has been introduced) in a suitable medium such that FCTR protein is produced. In another embodiment, the method further comprises isolating FCTR from the medium or the host cell.

Transgenic animals

The host cells of the invention can also be used to produce nonhuman transgenic animals. For example, in one embodiment, a host cell of the invention is a fertilized oocyte or an embryonic stem cell into which FCTR_X-coding sequences have been introduced. Such
5 host cells can then be used to create non-human transgenic animals in which exogenous FCTR_X sequences have been introduced into their genome or homologous recombinant animals in which endogenous FCTR_X sequences have been altered. Such animals are useful for studying the function and/or activity of FCTR_X and for identifying and/or evaluating modulators of FCTR_X activity. As used herein, a "transgenic animal" is a non-human animal,
10 preferably a mammal, more preferably a rodent such as a rat or mouse, in which one or more of the cells of the animal includes a transgene. Other examples of transgenic animals include non-human primates, sheep, dogs, cows, goats, chickens, amphibians, etc. A transgene is exogenous DNA that is integrated into the genome of a cell from which a transgenic animal develops and that remains in the genome of the mature animal, thereby directing the
15 expression of an encoded gene product in one or more cell types or tissues of the transgenic animal. As used herein, a "homologous recombinant animal" is a non-human animal, preferably a mammal, more preferably a mouse, in which an endogenous FCTR_X gene has been altered by homologous recombination between the endogenous gene and an exogenous DNA molecule introduced into a cell of the animal, *e.g.*, an embryonic cell of the animal, prior
20 to development of the animal.

A transgenic animal of the invention can be created by introducing FCTR_X-encoding nucleic acid into the male pronuclei of a fertilized oocyte, *e.g.*, by microinjection, retroviral infection, and allowing the oocyte to develop in a pseudopregnant female foster animal. The human FCTR_X cDNA sequence of SEQ ID NO:1, 2, 4, or 5 can be introduced as a transgene
25 into the genome of a non-human animal. Alternatively, a nonhuman homologue of the human FCTR_X gene, such as a mouse FCTR_X gene, can be isolated based on hybridization to the human FCTR_X cDNA (described further above) and used as a transgene. Intronic sequences and polyadenylation signals can also be included in the transgene to increase the efficiency of expression of the transgene. A tissue-specific regulatory sequence(s) can be operably linked
30 to the FCTR_X transgene to direct expression of FCTR_X protein to particular cells. Methods for generating transgenic animals via embryo manipulation and microinjection, particularly animals such as mice, have become conventional in the art and are described, for example, in U.S. Pat. Nos. 4,736,866; 4,870,009; and 4,873,191; and Hogan 1986, In: MANIPULATING THE MOUSE EMBRYO, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. Similar

methods are used for production of other transgenic animals. A transgenic founder animal can be identified based upon the presence of the FCTR_X transgene in its genome and/or expression of FCTR_X mRNA in tissues or cells of the animals. A transgenic founder animal can then be used to breed additional animals carrying the transgene. Moreover, transgenic
5 animals carrying a transgene encoding FCTR_X can further be bred to other transgenic animals carrying other transgenes.

To create a homologous recombinant animal, a vector is prepared which contains at least a portion of a FCTR_X gene into which a deletion, addition or substitution has been introduced to thereby alter, *e.g.*, functionally disrupt, the FCTR_X gene. The FCTR_X gene can
10 be a human gene (*e.g.*, the cDNA of SEQ ID NO:1, 2, 4, or 5) but more preferably, is a non-human homologue of a human FCTR_X gene. For example, a mouse homologue of human FCTR_X gene of SEQ ID NO:1, 2, 4, or 5 can be used to construct a homologous recombination vector suitable for altering an endogenous FCTR_X gene in the mouse genome. In one embodiment, the vector is designed such that, upon homologous recombination, the
15 endogenous FCTR_X gene is functionally disrupted (*i.e.*, no longer encodes a functional protein; also referred to as a "knock out" vector).

Alternatively, the vector can be designed such that, upon homologous recombination, the endogenous FCTR_X gene is mutated or otherwise altered but still encodes functional protein (*e.g.*, the upstream regulatory region can be altered to thereby alter the expression of
20 the endogenous FCTR_X protein). In the homologous recombination vector, the altered portion of the FCTR_X gene is flanked at its 5' and 3' ends by additional nucleic acid of the FCTR_X gene to allow for homologous recombination to occur between the exogenous FCTR_X gene carried by the vector and an endogenous FCTR_X gene in an embryonic stem cell. The additional flanking FCTR_X nucleic acid is of sufficient length for successful homologous
25 recombination with the endogenous gene. Typically, several kilobases of flanking DNA (both at the 5' and 3' ends) are included in the vector. See *e.g.*, Thomas *et al.* (1987) *Cell* 51:503 for a description of homologous recombination vectors. The vector is introduced into an embryonic stem cell line (*e.g.*, by electroporation) and cells in which the introduced FCTR_X gene has homologously recombined with the endogenous FCTR_X gene are selected (see *e.g.*,
30 Li *et al.* (1992) *Cell* 69:915).

The selected cells are then injected into a blastocyst of an animal (*e.g.*, a mouse) to form aggregation chimeras. See *e.g.*, Bradley 1987, In: TERATOCARCINOMAS AND EMBRYONIC STEM CELLS: A PRACTICAL APPROACH, Robertson, ed. IRL, Oxford, pp. 113-152. A chimeric embryo can then be implanted into a suitable pseudopregnant female foster animal and the

embryo brought to term. Progeny harboring the homologously recombined DNA in their germ cells can be used to breed animals in which all cells of the animal contain the homologously recombined DNA by germline transmission of the transgene. Methods for constructing homologous recombination vectors and homologous recombinant animals are described
5 further in Bradley (1991) *Curr Opin Biotechnol* 2:823-829; PCT International Publication Nos.: WO 90/11354; WO 91/01140; WO 92/0968; and WO 93/04169.

In another embodiment, transgenic non-human animals can be produced that contain selected systems that allow for regulated expression of the transgene. One example of such a system is the cre/loxP recombinase system of bacteriophage P1. For a description of the
10 cre/loxP recombinase system, see, e.g., Lakso *et al.* (1992) *PNAS* 89:6232-6236. Another example of a recombinase system is the FLP recombinase system of *Saccharomyces cerevisiae* (O'Gorman *et al.* (1991) *Science* 251:1351-1355. If a cre/loxP recombinase system is used to regulate expression of the transgene, animals containing transgenes encoding both the Cre recombinase and a selected protein are required. Such animals can be provided
15 through the construction of "double" transgenic animals, e.g., by mating two transgenic animals, one containing a transgene encoding a selected protein and the other containing a transgene encoding a recombinase.

Clones of the non-human transgenic animals described herein can also be produced according to the methods described in Wilmut *et al.* (1997) *Nature* 385:810-813. In brief, a
20 cell, e.g., a somatic cell, from the transgenic animal can be isolated and induced to exit the growth cycle and enter G₀ phase. The quiescent cell can then be fused, e.g., through the use of electrical pulses, to an enucleated oocyte from an animal of the same species from which the quiescent cell is isolated. The reconstructed oocyte is then cultured such that it develops to morula or blastocyte and then transferred to pseudopregnant female foster animal. The
25 offspring borne of this female foster animal will be a clone of the animal from which the cell, e.g., the somatic cell, is isolated.

Pharmaceutical Compositions

The FCTR_X nucleic acid molecules, FCTR_X proteins, and anti-FCTR_X antibodies
30 (also referred to herein as "active compounds") of the invention, and derivatives, fragments, analogs and homologs thereof, can be incorporated into pharmaceutical compositions suitable for administration. Such compositions typically comprise the nucleic acid molecule, protein, or antibody and a pharmaceutically acceptable carrier. As used herein, "pharmaceutically acceptable carrier" is intended to include any and all solvents, dispersion media, coatings,

antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the compositions is contemplated. Supplementary active compounds can also be incorporated into the compositions.

A pharmaceutical composition of the invention is formulated to be compatible with its intended route of administration. Examples of routes of administration include parenteral, *e.g.*, intravenous, intradermal, subcutaneous, oral (*e.g.*, inhalation), transdermal (topical), transmucosal, and rectal administration. Solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates, and agents for the adjustment of tonicity such as sodium chloride or dextrose. The pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. For intravenous administration, suitable carriers include physiological saline, bacteriostatic water, Cremophor EL™ (BASF, Parsippany, N.J.) or phosphate buffered saline (PBS). In all cases, the composition must be sterile and should be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents,

for example, sugars, polyalcohols such as manitol, sorbitol, sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate and gelatin.

5 Sterile injectable solutions can be prepared by incorporating the active compound (*e.g.*, an FCTR_X protein or anti-FCTR_X antibody) in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle that contains a basic dispersion medium and the required other ingredients from
10 those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, methods of preparation are vacuum drying and freeze-drying that yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

Oral compositions generally include an inert diluent or an edible carrier. They can be
15 enclosed in gelatin capsules or compressed into tablets. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and used in the form of tablets, troches, or capsules. Oral compositions can also be prepared using a fluid carrier for use as a mouthwash, wherein the compound in the fluid carrier is applied orally and swished and expectorated or swallowed. Pharmaceutically compatible binding agents, and/or
20 adjuvant materials can be included as part of the composition. The tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate or Sterotes; a glidant such as colloidal silicon dioxide; a
25 sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring.

For administration by inhalation, the compounds are delivered in the form of an aerosol spray from pressured container or dispenser which contains a suitable propellant, *e.g.*, a gas such as carbon dioxide, or a nebulizer.

30 Systemic administration can also be by transmucosal or transdermal means. For transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art, and include, for example, for transmucosal administration, detergents, bile salts, and fusidic acid derivatives. Transmucosal administration can be accomplished through the use of nasal sprays

or suppositories. For transdermal administration, the active compounds are formulated into ointments, salves, gels, or creams as generally known in the art.

The compounds can also be prepared in the form of suppositories (*e.g.*, with conventional suppository bases such as cocoa butter and other glycerides) or retention enemas for rectal
5 delivery.

In one embodiment, the active compounds are prepared with carriers that will protect the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides,
10 polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Methods for preparation of such formulations will be apparent to those skilled in the art. The materials can also be obtained commercially from Alza Corporation and Nova Pharmaceuticals, Inc. Liposomal suspensions (including liposomes targeted to infected cells with monoclonal antibodies to viral antigens) can also be used as pharmaceutically acceptable carriers. These can be prepared
15 according to methods known to those skilled in the art, for example, as described in U.S. Pat. No. 4,522,811.

It is especially advantageous to formulate oral or parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subject to be treated;
20 each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by and directly dependent on the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and the limitations inherent in the art of compounding such an active compound for
25 the treatment of individuals.

The nucleic acid molecules of the invention can be inserted into vectors and used as gene therapy vectors. Gene therapy vectors can be delivered to a subject by, for example, intravenous injection, local administration (see U.S. Pat. No. 5,328,470) or by stereotactic injection (see *e.g.*, Chen *et al.* (1994) *PNAS* 91:3054-3057). The pharmaceutical preparation
30 of the gene therapy vector can include the gene therapy vector in an acceptable diluent, or can comprise a slow release matrix in which the gene delivery vehicle is imbedded. Alternatively, where the complete gene delivery vector can be produced intact from recombinant cells, *e.g.*, retroviral vectors, the pharmaceutical preparation can include one or more cells that produce the gene delivery system.

The pharmaceutical compositions can be included in a container, pack, or dispenser together with instructions for administration.

Uses and Methods of the Invention

5 The isolated nucleic acid molecules of the invention can be used to express FCTR_X protein (*e.g.*, via a recombinant expression vector in a host cell in gene therapy applications), to detect FCTR_X mRNA (*e.g.*, in a biological sample) or a genetic lesion in an FCTR_X gene, and to modulate FCTR_X activity, as described further below. In addition, the FCTR_X proteins can be used to screen drugs or compounds that modulate the FCTR_X activity or expression as
10 well as to treat disorders characterized by insufficient or excessive production of FCTR_X protein or production of FCTR_X protein forms that have decreased or aberrant activity compared to FCTR_X wild type protein (*e.g.* proliferative disorders such as cancer or preclampsia, immune system disorders and inflammation, neurological disorders, and skin and muscle abnormalities). In addition, the anti-FCTR_X antibodies of the invention can be used to
15 detect and isolate FCTR_X proteins and modulate FCTR_X activity.

This invention further pertains to novel agents identified by the above described screening assays and uses thereof for treatments as described herein.

Screening Assays

20 The invention provides a method (also referred to herein as a "screening assay") for identifying modulators, *i.e.*, candidate or test compounds or agents (*e.g.*, peptides, peptidomimetics, small molecules or other drugs) that bind to FCTR_X proteins or have a stimulatory or inhibitory effect on, for example, FCTR_X expression or FCTR_X activity.

In one embodiment, the invention provides assays for screening candidate or test compounds which bind to or modulate the activity of a membrane-bound form of an FCTR_X
25 protein or polypeptide or biologically active portion thereof. The test compounds of the present invention can be obtained using any of the numerous approaches in combinatorial library methods known in the art, including: biological libraries; spatially addressable parallel solid phase or solution phase libraries; synthetic library methods requiring deconvolution; the
"one-bead one-compound" library method; and synthetic library methods using affinity
30 chromatography selection. The biological library approach is limited to peptide libraries, while the other four approaches are applicable to peptide, non-peptide oligomer or small molecule libraries of compounds (Lam (1997) *Anticancer Drug Des* 12:145).

Examples of methods for the synthesis of molecular libraries can be found in the art, for example in: DeWitt *et al.* (1993) *Proc Natl Acad Sci U.S.A.* 90:6909; Erb *et al.* (1994)

Proc Natl Acad Sci U.S.A. 91:11422; Zuckermann *et al.* (1994) *J Med Chem* 37:2678; Cho *et al.* (1993) *Science* 261:1303; Carrell *et al.* (1994) *Angew Chem Int Ed Engl* 33:2059; Carell *et al.* (1994) *Angew Chem Int Ed Engl* 33:2061; and Gallop *et al.* (1994) *J Med Chem* 37:1233.

Libraries of compounds may be presented in solution (*e.g.*, Houghten (1992)

- 5 *Biotechniques* 13:412-421), or on beads (Lam (1991) *Nature* 354:82-84), on chips (Fodor (1993) *Nature* 364:555-556), bacteria (Ladner U.S. Pat. No. 5,223,409), spores (Ladner USP '409), plasmids (Cull *et al.* (1992) *Proc Natl Acad Sci USA* 89:1865-1869) or on phage (Scott and Smith (1990) *Science* 249:386-390; Devlin (1990) *Science* 249:404-406; Cwirla *et al.* (1990) *Proc Natl Acad Sci U.S.A.* 87:6378-6382; Felici (1991) *J Mol Biol* 222:301-310; 10 Ladner above.).

- In one embodiment, an assay is a cell-based assay in which a cell which expresses a membrane-bound form of FCTR_X protein, or a biologically active portion thereof, on the cell surface is contacted with a test compound and the ability of the test compound to bind to an FCTR_X protein is determined. The cell, for example, can be of mammalian origin or a yeast 15 cell. Determining the ability of the test compound to bind to the FCTR_X protein can be accomplished, for example, by coupling the test compound with a radioisotope or enzymatic label such that binding of the test compound to the FCTR_X protein or biologically active portion thereof can be determined by detecting the labeled compound in a complex. For example, test compounds can be labeled with ¹²⁵I, ³⁵S, ¹⁴C, or ³H, either directly or indirectly, 20 and the radioisotope detected by direct counting of radioemission or by scintillation counting. Alternatively, test compounds can be enzymatically labeled with, for example, horseradish peroxidase, alkaline phosphatase, or luciferase, and the enzymatic label detected by determination of conversion of an appropriate substrate to product. In one embodiment, the assay comprises contacting a cell which expresses a membrane-bound form of FCTR_X 25 protein, or a biologically active portion thereof, on the cell surface with a known compound which binds FCTR_X to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with FCTR_X protein, wherein determining the ability of the test compound to interact with an FCTR_X protein comprises determining the ability of the test compound to preferentially bind to FCTR_X or a 30 biologically active portion thereof as compared to the known compound.

In another embodiment, an assay is a cell-based assay comprising contacting a cell expressing a membrane-bound form of FCTR_X protein, or a biologically active portion thereof, on the cell surface with a test compound and determining the ability of the test compound to modulate (*e.g.*, stimulate or inhibit) the activity of the FCTR_X protein or

biologically active portion thereof. Determining the ability of the test compound to modulate the activity of FCTR_X or a biologically active portion thereof can be accomplished, for example, by determining the ability of the FCTR_X protein to bind to or interact with an FCTR_X target molecule. As used herein, a "target molecule" is a molecule with which an FCTR_X protein binds or interacts in nature, for example, a molecule on the surface of a cell which expresses an FCTR_X protein, a molecule on the surface of a second cell, a molecule in the extracellular milieu, a molecule associated with the internal surface of a cell membrane or a cytoplasmic molecule. An FCTR_X target molecule can be a non-FCTR_X molecule or an FCTR_X protein or polypeptide of the present invention. In one embodiment, an FCTR_X target molecule is a component of a signal transduction pathway that facilitates transduction of an extracellular signal (*e.g.* a signal generated by binding of a compound to a membrane-bound FCTR_X molecule) through the cell membrane and into the cell. The target, for example, can be a second intercellular protein that has catalytic activity or a protein that facilitates the association of downstream signaling molecules with FCTR_X.

Determining the ability of the FCTR_X protein to bind to or interact with an FCTR_X target molecule can be accomplished by one of the methods described above for determining direct binding. In one embodiment, determining the ability of the FCTR_X protein to bind to or interact with an FCTR_X target molecule can be accomplished by determining the activity of the target molecule. For example, the activity of the target molecule can be determined by detecting induction of a cellular second messenger of the target (*i.e.* intracellular Ca^{2+} , diacylglycerol, IP_3 , etc.), detecting catalytic/enzymatic activity of the target an appropriate substrate, detecting the induction of a reporter gene (comprising an FCTR_X-responsive regulatory element operatively linked to a nucleic acid encoding a detectable marker, *e.g.*, luciferase), or detecting a cellular response, for example, cell survival, cellular differentiation, or cell proliferation.

In yet another embodiment, an assay of the present invention is a cell-free assay comprising contacting an FCTR_X protein or biologically active portion thereof with a test compound and determining the ability of the test compound to bind to the FCTR_X protein or biologically active portion thereof. Binding of the test compound to the FCTR_X protein can be determined either directly or indirectly as described above. In one embodiment, the assay comprises contacting the FCTR_X protein or biologically active portion thereof with a known compound which binds FCTR_X to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with an FCTR_X protein, wherein determining the ability of the test compound to interact with an FCTR_X

protein comprises determining the ability of the test compound to preferentially bind to FCTR_X or biologically active portion thereof as compared to the known compound.

5 In another embodiment, an assay is a cell-free assay comprising contacting FCTR_X protein or biologically active portion thereof with a test compound and determining the ability of the test compound to modulate (*e.g.* stimulate or inhibit) the activity of the FCTR_X protein or biologically active portion thereof. Determining the ability of the test compound to modulate the activity of FCTR_X can be accomplished, for example, by determining the ability of the FCTR_X protein to bind to an FCTR_X target molecule by one of the methods described above for determining direct binding. In an alternative embodiment, determining the ability of the test compound to modulate the activity of FCTR_X can be accomplished by determining the ability of the FCTR_X protein further modulate an FCTR_X target molecule. For example, the catalytic/enzymatic activity of the target molecule on an appropriate substrate can be determined as previously described.

10 In yet another embodiment, the cell-free assay comprises contacting the FCTR_X protein or biologically active portion thereof with a known compound which binds FCTR_X to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with an FCTR_X protein, wherein determining the ability of the test compound to interact with an FCTR_X protein comprises determining the ability of the FCTR_X protein to preferentially bind to or modulate the activity of an FCTR_X target molecule.

15 The cell-free assays of the present invention are amenable to use of both the soluble form or the membrane-bound form of FCTR_X. In the case of cell-free assays comprising the membrane-bound form of FCTR_X, it may be desirable to utilize a solubilizing agent such that the membrane-bound form of FCTR_X is maintained in solution. Examples of such solubilizing agents include non-ionic detergents such as n-octylglucoside, n-dodecylglucoside, n-dodecylmaltoside, octanoyl-N-methylglucamide, decanoyl-N-methylglucamide, Triton[®] X-100, Triton[®] X-114, Thesit[®], Isotridecypoly(ethylene glycol ether)_n, 3-(3-cholamidopropyl)dimethylamminiol-1-propane sulfonate (CHAPS), 3-(3-cholamidopropyl)dimethylamminiol-2-hydroxy-1-propane sulfonate (CHAPSO), or N-dodecyl--N,N-dimethyl-3-ammonio-1-propane sulfonate.

20 In more than one embodiment of the above assay methods of the present invention, it may be desirable to immobilize either FCTR_X or its target molecule to facilitate separation of complexed from uncomplexed forms of one or both of the proteins, as well as to accommodate automation of the assay. Binding of a test compound to FCTR_X, or interaction of FCTR_X

with a target molecule in the presence and absence of a candidate compound, can be accomplished in any vessel suitable for containing the reactants. Examples of such vessels include microtiter plates, test tubes, and micro-centrifuge tubes. In one embodiment, a fusion protein can be provided that adds a domain that allows one or both of the proteins to be bound to a matrix. For example, GST-FCTR_X fusion proteins or GST-target fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical, St. Louis, MO) or glutathione derivatized microtiter plates, that are then combined with the test compound or the test compound and either the non-adsorbed target protein or FCTR_X protein, and the mixture is incubated under conditions conducive to complex formation (*e.g.*, at physiological conditions for salt and pH). Following incubation, the beads or microtiter plate wells are washed to remove any unbound components, the matrix immobilized in the case of beads, complex determined either directly or indirectly, for example, as described above. Alternatively, the complexes can be dissociated from the matrix, and the level of FCTR_X binding or activity determined using standard techniques.

Other techniques for immobilizing proteins on matrices can also be used in the screening assays of the invention. For example, either FCTR_X or its target molecule can be immobilized utilizing conjugation of biotin and streptavidin. Biotinylated FCTR_X or target molecules can be prepared from biotin-NHS (N-hydroxy-succinimide) using techniques well known in the art (*e.g.*, biotinylation kit, Pierce Chemicals, Rockford, Ill.), and immobilized in the wells of streptavidin-coated 96 well plates (Pierce Chemical). Alternatively, antibodies reactive with FCTR_X or target molecules, but which do not interfere with binding of the FCTR_X protein to its target molecule, can be derivatized to the wells of the plate, and unbound target or FCTR_X trapped in the wells by antibody conjugation. Methods for detecting such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using antibodies reactive with the FCTR_X or target molecule, as well as enzyme-linked assays that rely on detecting an enzymatic activity associated with the FCTR_X or target molecule.

In another embodiment, modulators of FCTR_X expression are identified in a method wherein a cell is contacted with a candidate compound and the expression of FCTR_X mRNA or protein in the cell is determined. The level of expression of FCTR_X mRNA or protein in the presence of the candidate compound is compared to the level of expression of FCTR_X mRNA or protein in the absence of the candidate compound. The candidate compound can then be identified as a modulator of FCTR_X expression based on this comparison. For example, when expression of FCTR_X mRNA or protein is greater (statistically significantly

greater) in the presence of the candidate compound than in its absence, the candidate compound is identified as a stimulator of FCTR_X mRNA or protein expression. Alternatively, when expression of FCTR_X mRNA or protein is less (statistically significantly less) in the presence of the candidate compound than in its absence, the candidate compound is identified
5 as an inhibitor of FCTR_X mRNA or protein expression. The level of FCTR_X mRNA or protein expression in the cells can be determined by methods described herein for detecting FCTR_X mRNA or protein.

In yet another aspect of the invention, the FCTR_X proteins can be used as "bait proteins" in a two-hybrid assay or three hybrid assay (see, *e.g.*, U.S. Pat. No. 5,283,317;
10 Zervos *et al.* (1993) Cell 72:223-232; Madura *et al.* (1993) J Biol Chem 268:12046-12054; Bartel *et al.* (1993) Biotechniques 14:920-924; Iwabuchi *et al.* (1993) Oncogene 8:1693-1696; and Brent WO94/10300), to identify other proteins that bind to or interact with FCTR_X ("FCTR_X-binding proteins" or "FCTR_X-bps") and modulate FCTR_X activity. Such FCTR_X-binding proteins are also likely to be involved in the propagation of signals by the
15 FCTR_X proteins as, for example, upstream or downstream elements of the FCTR_X pathway.

The two-hybrid system is based on the modular nature of most transcription factors, which consist of separable DNA-binding and activation domains. Briefly, the assay utilizes two different DNA constructs. In one construct, the gene that codes for FCTR_X is fused to a gene encoding the DNA binding domain of a known transcription factor (*e.g.*, GAL-4). In the
20 other construct, a DNA sequence, from a library of DNA sequences, that encodes an unidentified protein ("prey" or "sample") is fused to a gene that codes for the activation domain of the known transcription factor. If the "bait" and the "prey" proteins are able to interact, *in vivo*, forming a FCTR_X-dependent complex, the DNA-binding and activation domains of the transcription factor are brought into close FCTR_Ximity. This FCTR_Ximity
25 allows transcription of a reporter gene (*e.g.*, LacZ) that is operably linked to a transcriptional regulatory site responsive to the transcription factor. Expression of the reporter gene can be detected and cell colonies containing the functional transcription factor can be isolated and used to obtain the cloned gene that encodes the protein which interacts with FCTR_X.

This invention further pertains to novel agents identified by the above-described
30 screening assays and uses thereof for treatments as described herein.

Detection Assays

Portions or fragments of the cDNA sequences identified herein (and the corresponding complete gene sequences) can be used in numerous ways as polynucleotide reagents. For

example, these sequences can be used to: (i) map their respective genes on a chromosome; and, thus, locate gene regions associated with genetic disease; (ii) identify an individual from a minute biological sample (tissue typing); and (iii) aid in forensic identification of a biological sample. These applications are described in the subsections below.

5

Chromosome Mapping

Once the sequence (or a portion of the sequence) of a gene has been isolated, this sequence can be used to map the location of the gene on a chromosome. This process is called chromosome mapping. Accordingly, portions or fragments of the FCTR_X sequences,
10 described herein, can be used to map the location of the FCTR_X genes, respectively, on a chromosome. The mapping of the FCTR_X sequences to chromosomes is an important first step in correlating these sequences with genes associated with disease.

Briefly, FCTR_X genes can be mapped to chromosomes by preparing PCR primers (preferably 15-25 bp in length) from the FCTR_X sequences. Computer analysis of the
15 FCTR_X sequences can be used to rapidly select primers that do not span more than one exon in the genomic DNA, thus complicating the amplification process. These primers can then be used for PCR screening of somatic cell hybrids containing individual human chromosomes. Only those hybrids containing the human gene corresponding to the FCTR_X sequences will yield an amplified fragment.

20 Somatic cell hybrids are prepared by fusing somatic cells from different mammals (e.g., human and mouse cells). As hybrids of human and mouse cells grow and divide, they gradually lose human chromosomes in random order, but retain the mouse chromosomes. By using media in which mouse cells cannot grow, because they lack a particular enzyme, but in which human cells can, the one human chromosome that contains the gene encoding the
25 needed enzyme will be retained. By using various media, panels of hybrid cell lines can be established. Each cell line in a panel contains either a single human chromosome or a small number of human chromosomes, and a full set of mouse chromosomes, allowing easy mapping of individual genes to specific human chromosomes. (D'Eustachio *et al.* (1983) *Science* 220:919-924). Somatic cell hybrids containing only fragments of human
30 chromosomes can also be produced by using human chromosomes with translocations and deletions.

PCR mapping of somatic cell hybrids is a rapid procedure for assigning a particular sequence to a particular chromosome. Three or more sequences can be assigned per day using

a single thermal cycler. Using the FCTR_X sequences to design oligonucleotide primers, sublocalization can be achieved with panels of fragments from specific chromosomes.

Fluorescence *in situ* hybridization (FISH) of a DNA sequence to a metaphase chromosomal spread can further be used to provide a precise chromosomal location in one
5 step. Chromosome spreads can be made using cells whose division has been blocked in metaphase by a chemical like colcemid that disrupts the mitotic spindle. The chromosomes can be treated briefly with trypsin, and then stained with Giemsa. A pattern of light and dark bands develops on each chromosome, so that the chromosomes can be identified individually. The FISH technique can be used with a DNA sequence as short as 500 or 600 bases.
10 However, clones larger than 1,000 bases have a higher likelihood of binding to a unique chromosomal location with sufficient signal intensity for simple detection. Preferably 1,000 bases, and more preferably 2,000 bases, will suffice to get good results at a reasonable amount of time. For a review of this technique, see Verma *et al.*, HUMAN CHROMOSOMES: A MANUAL OF BASIC TECHNIQUES (Pergamon Press, New York 1988).

15 Reagents for chromosome mapping can be used individually to mark a single chromosome or a single site on that chromosome, or panels of reagents can be used for marking multiple sites and/or multiple chromosomes. Reagents corresponding to noncoding regions of the genes actually are preferred for mapping purposes. Coding sequences are more likely to be conserved within gene families, thus increasing the chance of cross hybridizations
20 during chromosomal mapping.

Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. Such data are found, for example, in McKusick, MENDELIAN INHERITANCE IN MAN, available on-line through Johns Hopkins University Welch Medical Library). The relationship between
25 genes and disease, mapped to the same chromosomal region, can then be identified through linkage analysis (co-inheritance of physically adjacent genes), described in, for example, Egeland *et al.* (1987) *Nature*, 325:783-787.

Moreover, differences in the DNA sequences between individuals affected and unaffected with a disease associated with the FCTR_X gene, can be determined. If a mutation
30 is observed in some or all of the affected individuals but not in any unaffected individuals, then the mutation is likely to be the causative agent of the particular disease. Comparison of affected and unaffected individuals generally involves first looking for structural alterations in the chromosomes, such as deletions or translocations that are visible from chromosome spreads or detectable using PCR based on that DNA sequence. Ultimately, complete

sequencing of genes from several individuals can be performed to confirm the presence of a mutation and to distinguish mutations from polymorphisms.

Tissue Typing

5 The FCTR_X sequences of the present invention can also be used to identify individuals from minute biological samples. In this technique, an individual's genomic DNA is digested with one or more restriction enzymes, and probed on a Southern blot to yield unique bands for identification. The sequences of the present invention are useful as additional DNA markers for RFLP ("restriction fragment length polymorphisms," described in U.S. Pat. No.
10 5,272,057).

 Furthermore, the sequences of the present invention can be used to provide an alternative technique that determines the actual base-by-base DNA sequence of selected portions of an individual's genome. Thus, the FCTR_X sequences described herein can be used to prepare two PCR primers from the 5' and 3' ends of the sequences. These primers can then
15 be used to amplify an individual's DNA and subsequently sequence it.

 Panels of corresponding DNA sequences from individuals, prepared in this manner, can provide unique individual identifications, as each individual will have a unique set of such DNA sequences due to allelic differences. The sequences of the present invention can be used to obtain such identification sequences from individuals and from tissue. The FCTR_X
20 sequences of the invention uniquely represent portions of the human genome. Allelic variation occurs to some degree in the coding regions of these sequences, and to a greater degree in the noncoding regions. It is estimated that allelic variation between individual humans occurs with a frequency of about once per each 500 bases. Much of the allelic variation is due to single nucleotide polymorphisms (SNPs), which include restriction
25 fragment length polymorphisms (RFLPs).

 Each of the sequences described herein can, to some degree, be used as a standard against which DNA from an individual can be compared for identification purposes. Because greater numbers of polymorphisms occur in the noncoding regions, fewer sequences are necessary to differentiate individuals. For example, the noncoding sequences of SEQ ID
30 NO:1, 2, 4, or 5 can comfortably provide positive individual identification with a panel of perhaps 10 to 1,000 primers that each yield a noncoding amplified sequence of 100 bases. If predicted coding sequences, such as the nucleic acid sequences that code for the amino acid sequences of SEQ ID NO:3 or 6 are used, a more appropriate number of primers for positive individual identification would be 500-2,000.

Predictive Medicine

The present invention also pertains to the field of predictive medicine in which diagnostic assays, prognostic assays, pharmacogenomics, and monitoring clinical trials are used for prognostic (predictive) purposes to thereby treat an individual prophylactically. Accordingly, one aspect of the present invention relates to diagnostic assays for determining FCTR_X protein and/or nucleic acid expression as well as FCTR_X activity, in the context of a biological sample (*e.g.*, blood, serum, cells, tissue) to thereby determine whether an individual is afflicted with a disease or disorder, or is at risk of developing a disorder, associated with aberrant FCTR_X expression or activity. The invention also provides for prognostic (or predictive) assays for determining whether an individual is at risk of developing a disorder associated with FCTR_X protein, nucleic acid expression or activity. For example, mutations in an FCTR_X gene can be assayed in a biological sample. Such assays can be used for prognostic or predictive purpose to thereby prophylactically treat an individual prior to the onset of a disorder characterized by or associated with FCTR_X protein, nucleic acid expression or activity.

Another aspect of the invention provides methods for determining FCTR_X protein, nucleic acid expression or FCTR_X activity in an individual to thereby select appropriate therapeutic or prophylactic agents for that individual (referred to herein as "pharmacogenomics"). Pharmacogenomics allows for the selection of agents (*e.g.*, drugs) for therapeutic or prophylactic treatment of an individual based on the genotype of the individual (*e.g.*, the genotype of the individual examined to determine the ability of the individual to respond to a particular agent).

Yet another aspect of the invention pertains to monitoring the influence of agents (*e.g.*, drugs, compounds) on the expression or activity of FCTR_X in clinical trials.

These and other agents are described in further detail in the following sections.

Diagnostic Assays

An exemplary method for detecting the presence or absence of FCTR_X in a biological sample involves obtaining a biological sample from a test subject and contacting the biological sample with a compound or an agent capable of detecting FCTR_X protein or nucleic acid (*e.g.*, mRNA, genomic DNA) that encodes FCTR_X protein such that the presence of FCTR_X is detected in the biological sample. An agent for detecting FCTR_X mRNA or genomic DNA is a labeled nucleic acid probe capable of hybridizing to FCTR_X mRNA or genomic DNA. The

nucleic acid probe can be, for example, a full-length FCTR_X nucleic acid, such as the nucleic acid of SEQ ID NO:1, 2, 4, or 5, or a portion thereof, such as an oligonucleotide of at least 15, 30, 50, 100, 250 or 500 nucleotides in length and sufficient to specifically hybridize under stringent conditions to FCTR_X mRNA or genomic DNA. Other suitable probes for use in the
5 diagnostic assays of the invention are described herein.

An agent for detecting FCTR_X protein is an antibody capable of binding to FCTR_X protein, preferably an antibody with a detectable label. Antibodies can be polyclonal, or more preferably, monoclonal. An intact antibody, or a fragment thereof (*e.g.*, Fab or F(ab')₂) can be used. The term "labeled", with regard to the probe or antibody, is intended to encompass
10 direct labeling of the probe or antibody by coupling (*i.e.*, physically linking) a detectable substance to the probe or antibody, as well as indirect labeling of the probe or antibody by reactivity with another reagent that is directly labeled. Examples of indirect labeling include detection of a primary antibody using a fluorescently labeled secondary antibody and end-labeling of a DNA probe with biotin such that it can be detected with fluorescently labeled
15 streptavidin. The term "biological sample" is intended to include tissues, cells and biological fluids isolated from a subject, as well as tissues, cells and fluids present within a subject. That is, the detection method of the invention can be used to detect FCTR_X mRNA, protein, or genomic DNA in a biological sample *in vitro* as well as *in vivo*. For example, *in vitro* techniques for detection of FCTR_X mRNA include Northern hybridizations and *in situ*
20 hybridizations. *In vitro* techniques for detection of FCTR_X protein include enzyme linked immunosorbent assays (ELISAs), Western blots, immunoprecipitations and immunofluorescence. *In vitro* techniques for detection of FCTR_X genomic DNA include Southern hybridizations. Furthermore, *in vivo* techniques for detection of FCTR_X protein include introducing into a subject a labeled anti-FCTR_X antibody. For example, the antibody
25 can be labeled with a radioactive marker whose presence and location in a subject can be detected by standard imaging techniques.

In one embodiment, the biological sample contains protein molecules from the test subject. Alternatively, the biological sample can contain mRNA molecules from the test subject or genomic DNA molecules from the test subject. A preferred biological sample is a
30 peripheral blood leukocyte sample isolated by conventional means from a subject.

In another embodiment, the methods further involve obtaining a control biological sample from a control subject, contacting the control sample with a compound or agent capable of detecting FCTR_X protein, mRNA, or genomic DNA, such that the presence of FCTR_X protein, mRNA or genomic DNA is detected in the biological sample, and comparing

the presence of FCTR_X protein, mRNA or genomic DNA in the control sample with the presence of FCTR_X protein, mRNA or genomic DNA in the test sample.

The invention also encompasses kits for detecting the presence of FCTR_X in a biological sample. For example, the kit can comprise: a labeled compound or agent capable of
5 detecting FCTR_X protein or mRNA in a biological sample; means for determining the amount of FCTR_X in the sample; and means for comparing the amount of FCTR_X in the sample with a standard. The compound or agent can be packaged in a suitable container. The kit can further comprise instructions for using the kit to detect FCTR_X protein or nucleic acid.

10 **Prognostic Assays**

The diagnostic methods described herein can furthermore be utilized to identify subjects having or at risk of developing a disease or disorder associated with aberrant FCTR_X expression or activity. For example, the assays described herein, such as the preceding diagnostic assays or the following assays, can be utilized to identify a subject having or at risk
15 of developing a disorder associated with FCTR_X protein, nucleic acid expression or activity such as cancer or fibrotic disorders. Alternatively, the prognostic assays can be utilized to identify a subject having or at risk for developing a disease or disorder. Thus, the present invention provides a method for identifying a disease or disorder associated with aberrant FCTR_X expression or activity in which a test sample is obtained from a subject and FCTR_X
20 protein or nucleic acid (*e.g.*, mRNA, genomic DNA) is detected, wherein the presence of FCTR_X protein or nucleic acid is diagnostic for a subject having or at risk of developing a disease or disorder associated with aberrant FCTR_X expression or activity. As used herein, a "test sample" refers to a biological sample obtained from a subject of interest. For example, a test sample can be a biological fluid (*e.g.*, serum), cell sample, or tissue.

25 Furthermore, the prognostic assays described herein can be used to determine whether a subject can be administered an agent (*e.g.*, an agonist, antagonist, peptidomimetic, protein, peptide, nucleic acid, small molecule, or other drug candidate) to treat a disease or disorder associated with aberrant FCTR_X expression or activity. For example, such methods can be used to determine whether a subject can be effectively treated with an agent for a disorder,
30 such as cancer or preclampsia. Thus, the present invention provides methods for determining whether a subject can be effectively treated with an agent for a disorder associated with aberrant FCTR_X expression or activity in which a test sample is obtained and FCTR_X protein or nucleic acid is detected (*e.g.*, wherein the presence of FCTR_X protein or nucleic acid is

diagnostic for a subject that can be administered the agent to treat a disorder associated with aberrant FCTR_X expression or activity.)

The methods of the invention can also be used to detect genetic lesions in an FCTR_X gene, thereby determining if a subject with the lesioned gene is at risk for a disorder characterized by aberrant cell proliferation and/or differentiation. In various embodiments, the methods include detecting, in a sample of cells from the subject, the presence or absence of a genetic lesion characterized by at least one of an alteration affecting the integrity of a gene encoding an FCTR_X-protein, or the mis-expression of the FCTR_X gene. For example, such genetic lesions can be detected by ascertaining the existence of at least one of (1) a deletion of one or more nucleotides from an FCTR_X gene; (2) an addition of one or more nucleotides to an FCTR_X gene; (3) a substitution of one or more nucleotides of an FCTR_X gene, (4) a chromosomal rearrangement of an FCTR_X gene; (5) an alteration in the level of a messenger RNA transcript of an FCTR_X gene, (6) aberrant modification of an FCTR_X gene, such as of the methylation pattern of the genomic DNA, (7) the presence of a non-wild type splicing pattern of a messenger RNA transcript of an FCTR_X gene, (8) a non-wild type level of an FCTR_X-protein, (9) allelic loss of an FCTR_X gene, and (10) inappropriate post-translational modification of an FCTR_X-protein. As described herein, there are a large number of assay techniques known in the art which can be used for detecting lesions in an FCTR_X gene. A preferred biological sample is a peripheral blood leukocyte sample isolated by conventional means from a subject. However, any biological sample containing nucleated cells may be used, including, for example, buccal mucosal cells.

In certain embodiments, detection of the lesion involves the use of a probe/primer in a polymerase chain reaction (PCR) (see, *e.g.*, U.S. Pat. Nos. 4,683,195 and 4,683,202), such as anchor PCR or RACE PCR, or, alternatively, in a ligation chain reaction (LCR) (see, *e.g.*, Landegran *et al.* (1988) *Science* 241:1077-1080; and Nakazawa *et al.* (1994) *PNAS* 91:360-364), the latter of which can be particularly useful for detecting point mutations in an FCTR_X-gene (see Abravaya *et al.* (1995) *Nucl Acids Res* 23:675-682). This method can include the steps of collecting a sample of cells from a patient, isolating nucleic acid (*e.g.*, genomic, mRNA or both) from the cells of the sample, contacting the nucleic acid sample with one or more primers that specifically hybridize to an FCTR_X gene under conditions such that hybridization and amplification of the FCTR_X gene (if present) occurs, and detecting the presence or absence of an amplification product, or detecting the size of the amplification product and comparing the length to a control sample. It is anticipated that PCR and/or LCR

may be desirable to use as a preliminary amplification step in conjunction with any of the techniques used for detecting mutations described herein.

Alternative amplification methods include: self sustained sequence replication (Guatelli *et al.*, 1990, *Proc Natl Acad Sci USA* 87:1874-1878), transcriptional amplification
5 system (Kwoh, *et al.*, 1989, *Proc Natl Acad Sci USA* 86:1173-1177), Q-Beta Replicase (Lizardi *et al.*, 1988, *BioTechnology* 6:1197), or any other nucleic acid amplification method, followed by the detection of the amplified molecules using techniques well known to those of skill in the art. These detection schemes are especially useful for the detection of nucleic acid molecules if such molecules are present in very low numbers.

10 In an alternative embodiment, mutations in an FCTRX gene from a sample cell can be identified by alterations in restriction enzyme cleavage patterns. For example, sample and control DNA is isolated, amplified (optionally), digested with one or more restriction endonucleases, and fragment length sizes are determined by gel electrophoresis and compared. Differences in fragment length sizes between sample and control DNA indicate mutations in
15 the sample DNA. Moreover, the use of sequence specific ribozymes (see, for example, U.S. Pat. No. 5,493,531) can be used to score for the presence of specific mutations by development or loss of a ribozyme cleavage site.

In other embodiments, genetic mutations in FCTRX can be identified by hybridizing a sample and control nucleic acids, *e.g.*, DNA or RNA, to high density arrays containing
20 hundreds or thousands of oligonucleotides probes (Cronin *et al.* (1996) *Human Mutation* 7: 244-255; Kozal *et al.* (1996) *Nature Medicine* 2: 753-759). For example, genetic mutations in FCTRX can be identified in two dimensional arrays containing light-generated DNA probes as described in Cronin *et al.* above. Briefly, a first hybridization array of probes can be used to scan through long stretches of DNA in a sample and control to identify base changes between
25 the sequences by making linear arrays of sequential overlapping probes. This step allows the identification of point mutations. This step is followed by a second hybridization array that allows the characterization of specific mutations by using smaller, specialized probe arrays complementary to all variants or mutations detected. Each mutation array is composed of parallel probe sets, one complementary to the wild-type gene and the other complementary to
30 the mutant gene.

In yet another embodiment, any of a variety of sequencing reactions known in the art can be used to directly sequence the FCTRX gene and detect mutations by comparing the sequence of the sample FCTRX with the corresponding wild-type (control) sequence. Examples of sequencing reactions include those based on techniques developed by Maxim and

Gilbert (1977) *PNAS* 74:560 or Sanger (1977) *PNAS* 74:5463. It is also contemplated that any of a variety of automated sequencing procedures can be utilized when performing the diagnostic assays (Naeve *et al.*, (1995) *Biotechniques* 19:448), including sequencing by mass spectrometry (see, *e.g.*, PCT International Publ. No. WO 94/16101; Cohen *et al.* (1996) *Adv Chromatogr* 36:127-162; and Griffin *et al.* (1993) *Appl Biochem Biotechnol* 38:147-159).

Other methods for detecting mutations in the FCTR_X gene include methods in which protection from cleavage agents is used to detect mismatched bases in RNA/RNA or RNA/DNA heteroduplexes (Myers *et al.* (1985) *Science* 230:1242). In general, the art technique of "mismatch cleavage" starts by providing heteroduplexes of formed by hybridizing (labeled) RNA or DNA containing the wild-type FCTR_X sequence with potentially mutant RNA or DNA obtained from a tissue sample. The double-stranded duplexes are treated with an agent that cleaves single-stranded regions of the duplex such as which will exist due to basepair mismatches between the control and sample strands. For instance, RNA/DNA duplexes can be treated with RNase and DNA/DNA hybrids treated with S1 nuclease to enzymatically digesting the mismatched regions. In other embodiments, either DNA/DNA or RNA/DNA duplexes can be treated with hydroxylamine or osmium tetroxide and with piperidine in order to digest mismatched regions. After digestion of the mismatched regions, the resulting material is then separated by size on denaturing polyacrylamide gels to determine the site of mutation. See, for example, Cotton *et al* (1988) *Proc Natl Acad Sci USA* 85:4397; Saleeba *et al* (1992) *Methods Enzymol* 217:286-295. In an embodiment, the control DNA or RNA can be labeled for detection.

In still another embodiment, the mismatch cleavage reaction employs one or more proteins that recognize mismatched base pairs in double-stranded DNA (so called "DNA mismatch repair" enzymes) in defined systems for detecting and mapping point mutations in FCTR_X cDNAs obtained from samples of cells. For example, the mutY enzyme of *E. coli* cleaves A at G/A mismatches and the thymidine DNA glycosylase from HeLa cells cleaves T at G/T mismatches (Hsu *et al.* (1994) *Carcinogenesis* 15:1657-1662). According to an exemplary embodiment, a probe based on an FCTR_X sequence, *e.g.*, a wild-type FCTR_X sequence, is hybridized to a cDNA or other DNA product from a test cell(s). The duplex is treated with a DNA mismatch repair enzyme, and the cleavage products, if any, can be detected from electrophoresis protocols or the like. See, for example, U.S. Pat. No. 5,459,039.

In other embodiments, alterations in electrophoretic mobility will be used to identify mutations in FCTR_X genes. For example, single strand conformation polymorphism (SSCP) may be used to detect differences in electrophoretic mobility between mutant and wild type

nucleic acids (Orita *et al.* (1989) *Proc Natl Acad Sci USA*: 86:2766, see also Cotton (1993) *Mutat Res* 285:125-144; Hayashi (1992) *Genet Anal Tech Appl* 9:73-79). Single-stranded DNA fragments of sample and control FCTR_X nucleic acids will be denatured and allowed to renature. The secondary structure of single-stranded nucleic acids varies according to
5 sequence, the resulting alteration in electrophoretic mobility enables the detection of even a single base change. The DNA fragments may be labeled or detected with labeled probes. The sensitivity of the assay may be enhanced by using RNA (rather than DNA), in which the secondary structure is more sensitive to a change in sequence. In one embodiment, the subject method utilizes heteroduplex analysis to separate double stranded heteroduplex molecules on
10 the basis of changes in electrophoretic mobility (Keen *et al.* (1991) *Trends Genet* 7:5).

In yet another embodiment the movement of mutant or wild-type fragments in polyacrylamide gels containing a gradient of denaturant is assayed using denaturing gradient gel electrophoresis (DGGE) (Myers *et al.* (1985) *Nature* 313:495). When DGGE is used as the method of analysis, DNA will be modified to insure that it does not completely denature, for
15 example by adding a GC clamp of approximately 40 bp of high-melting GC-rich DNA by PCR. In a further embodiment, a temperature gradient is used in place of a denaturing gradient to identify differences in the mobility of control and sample DNA (Rosenbaum and Reissner (1987) *Biophys Chem* 265:12753).

Examples of other techniques for detecting point mutations include, but are not limited
20 to, selective oligonucleotide hybridization, selective amplification, or selective primer extension. For example, oligonucleotide primers may be prepared in which the known mutation is placed centrally and then hybridized to target DNA under conditions that permit hybridization only if a perfect match is found (Saiki *et al.* (1986) *Nature* 324:163); Saiki *et al.* (1989) *Proc Natl Acad. Sci USA* 86:6230). Such allele specific oligonucleotides are
25 hybridized to PCR amplified target DNA or a number of different mutations when the oligonucleotides are attached to the hybridizing membrane and hybridized with labeled target DNA.

Alternatively, allele specific amplification technology that depends on selective PCR amplification may be used in conjunction with the instant invention. Oligonucleotides used as
30 primers for specific amplification may carry the mutation of interest in the center of the molecule (so that amplification depends on differential hybridization) (Gibbs *et al.* (1989) *Nucleic Acids Res* 17:2437-2448) or at the extreme 3' end of one primer where, under appropriate conditions, mismatch can prevent, or reduce polymerase extension (Prossner (1993) *Tibtech* 11:238). In addition it may be desirable to introduce a novel restriction site in

the region of the mutation to create cleavage-based detection (Gasparini *et al* (1992) *Mol Cell Probes* 6:1). It is anticipated that in certain embodiments amplification may also be performed using Taq ligase for amplification (Barany (1991) *Proc Natl Acad Sci USA* 88:189). In such cases, ligation will occur only if there is a perfect match at the 3' end of the 5' sequence, making it possible to detect the presence of a known mutation at a specific site by looking for the presence or absence of amplification.

The methods described herein may be performed, for example, by utilizing pre-packaged diagnostic kits comprising at least one probe nucleic acid or antibody reagent described herein, which may be conveniently used, *e.g.*, in clinical settings to diagnose patients exhibiting symptoms or family history of a disease or illness involving an FCTR_X gene.

Furthermore, any cell type or tissue, preferably peripheral blood leukocytes, in which FCTR_X is expressed may be utilized in the prognostic assays described herein. However, any biological sample containing nucleated cells may be used, including, for example, buccal mucosal cells.

Pharmacogenomics

Agents, or modulators that have a stimulatory or inhibitory effect on FCTR_X activity (*e.g.*, FCTR_X gene expression), as identified by a screening assay described herein can be administered to individuals to treat (prophylactically or therapeutically) disorders (*e.g.*, cancer or immune disorders, neurological disorders, muscular dystrophy, or epidermolysis bullosa simplex) associated with aberrant FCTR_X activity. In conjunction with such treatment, the pharmacogenomics (*i.e.*, the study of the relationship between an individual's genotype and that individual's response to a foreign compound or drug) of the individual may be considered. Differences in metabolism of therapeutics can lead to severe toxicity or therapeutic failure by altering the relation between dose and blood concentration of the pharmacologically active drug. Thus, the pharmacogenomics of the individual permits the selection of effective agents (*e.g.*, drugs) for prophylactic or therapeutic treatments based on a consideration of the individual's genotype. Such pharmacogenomics can further be used to determine appropriate dosages and therapeutic regimens. Accordingly, the activity of FCTR_X protein, expression of FCTR_X nucleic acid, or mutation content of FCTR_X genes in an individual can be determined to thereby select appropriate agent(s) for therapeutic or prophylactic treatment of the individual.

Pharmacogenomics deals with clinically significant hereditary variations in the response to drugs due to altered drug disposition and abnormal action in affected persons. See *e.g.*, Eichelbaum, *Clin Exp Pharmacol Physiol*, 1996, 23:983-985 and Linder, *Clin Chem*, 1997, 43:254-266. In general, two types of pharmacogenetic conditions can be differentiated.

- 5 Genetic conditions transmitted as a single factor altering the way drugs act on the body (altered drug action) or genetic conditions transmitted as single factors altering the way the body acts on drugs (altered drug metabolism). These pharmacogenetic conditions can occur either as rare defects or as polymorphisms. For example, glucose-6-phosphate dehydrogenase (G6PD) deficiency is a common inherited enzymopathy in which the main clinical
10 complication is haemolysis after ingestion of oxidant drugs (anti-malarials, sulfonamides, analgesics, nitrofurans) and consumption of fava beans.

- As an illustrative embodiment, the activity of drug metabolizing enzymes is a major determinant of both the intensity and duration of drug action. The discovery of genetic polymorphisms of drug metabolizing enzymes (*e.g.*, N-acetyltransferase 2 (NAT 2) and
15 cytochrome P450 enzymes CYP2D6 and CYP2C19) has provided an explanation as to why some patients do not obtain the expected drug effects or show exaggerated drug response and serious toxicity after taking the standard and safe dose of a drug. These polymorphisms are expressed in two phenotypes in the population, the extensive metabolizer (EM) and poor metabolizer (PM). The prevalence of PM is different among different populations. For
20 example, the gene coding for CYP2D6 is highly polymorphic and several mutations have been identified in PM, which all lead to the absence of functional CYP2D6. Poor metabolizers of CYP2D6 and CYP2C19 quite frequently experience exaggerated drug response and side effects when they receive standard doses. If a metabolite is the active therapeutic moiety, PM show no therapeutic response, as demonstrated for the analgesic effect of codeine mediated by
25 its CYP2D6-formed metabolite morphine. The other extreme is the so called ultra-rapid metabolizers who do not respond to standard doses. Recently, the molecular basis of ultra-rapid metabolism has been identified to be due to CYP2D6 gene amplification.

- Thus, the activity of FCTR protein, expression of FCTR nucleic acid, or mutation content of FCTR genes in an individual can be determined to thereby select appropriate
30 agent(s) for therapeutic or prophylactic treatment of the individual. In addition, pharmacogenetic studies can be used to apply genotyping of polymorphic alleles encoding drug-metabolizing enzymes to the identification of an individual's drug responsiveness phenotype. This knowledge, when applied to dosing or drug selection, can avoid adverse reactions or therapeutic failure and thus enhance therapeutic or prophylactic efficiency when

treating a subject with an FCTR_X modulator, such as a modulator identified by one of the exemplary screening assays described herein.

Monitoring of Effects During Clinical Trials

5 Monitoring the influence of agents (*e.g.*, drugs, compounds) on the expression or activity of FCTR_X (*e.g.*, the ability to modulate aberrant cell proliferation and/or differentiation) can be applied not only in basic drug screening, but also in clinical trials. For example, the effectiveness of an agent determined by a screening assay as described herein to increase FCTR_X gene expression, protein levels, or upregulate FCTR_X activity, can be
10 monitored in clinical trials of subjects exhibiting decreased FCTR_X gene expression, protein levels, or downregulated FCTR_X activity. Alternatively, the effectiveness of an agent determined by a screening assay to decrease FCTR_X gene expression, protein levels, or downregulate FCTR_X activity, can be monitored in clinical trials of subjects exhibiting increased FCTR_X gene expression, protein levels, or upregulated FCTR_X activity. In such
15 clinical trials, the expression or activity of FCTR_X and, preferably, other genes that have been implicated in, for example, a cellular proliferation disorder can be used as a "read out" or markers of the immune responsiveness of a particular cell.

For example, and not by way of limitation, genes, including FCTR_X, that are modulated in cells by treatment with an agent (*e.g.*, compound, drug or small molecule) that
20 modulates FCTR_X activity (*e.g.*, identified in a screening assay as described herein) can be identified. Thus, to study the effect of agents on cellular proliferation disorders, for example, in a clinical trial, cells can be isolated and RNA prepared and analyzed for the levels of expression of FCTR_X and other genes implicated in the disorder. The levels of gene expression (*i.e.*, a gene expression pattern) can be quantified by Northern blot analysis or
25 RT-PCR, as described herein, or alternatively by measuring the amount of protein produced, by one of the methods as described herein, or by measuring the levels of activity of FCTR_X or other genes. In this way, the gene expression pattern can serve as a marker, indicative of the physiological response of the cells to the agent. Accordingly, this response state may be determined before, and at various points during, treatment of the individual with the agent.

30 In one embodiment, the present invention provides a method for monitoring the effectiveness of treatment of a subject with an agent (*e.g.*, an agonist, antagonist, protein, peptide, peptidomimetic, nucleic acid, small molecule, or other drug candidate identified by the screening assays described herein) comprising the steps of (i) obtaining a pre-administration sample from a subject prior to administration of the agent; (ii) detecting the

level of expression of an FCTR_X protein, mRNA, or genomic DNA in the preadministration sample; (iii) obtaining one or more post-administration samples from the subject; (iv) detecting the level of expression or activity of the FCTR_X protein, mRNA, or genomic DNA in the post-administration samples; (v) comparing the level of expression or activity of the FCTR_X protein, mRNA, or genomic DNA in the pre-administration sample with the FCTR_X protein, mRNA, or genomic DNA in the post administration sample or samples; and (vi) altering the administration of the agent to the subject accordingly. For example, increased administration of the agent may be desirable to increase the expression or activity of FCTR_X to higher levels than detected, *i.e.*, to increase the effectiveness of the agent. Alternatively, decreased administration of the agent may be desirable to decrease expression or activity of FCTR_X to lower levels than detected, *i.e.*, to decrease the effectiveness of the agent.

Methods of Treatment

The present invention provides for both prophylactic and therapeutic methods of treating a subject at risk of (or susceptible to) a disorder or having a disorder associated with aberrant FCTR_X expression or activity.

Disorders

Diseases and disorders that are characterized by increased (relative to a subject not suffering from the disease or disorder) levels or biological activity may be treated with Therapeutics that antagonize (*i.e.*, reduce or inhibit) activity. Therapeutics that antagonize activity may be administered in a therapeutic or prophylactic manner. Therapeutics that may be utilized include, but are not limited to, (i) an aforementioned peptide, or analogs, derivatives, fragments or homologs thereof; (ii) antibodies to an aforementioned peptide; (iii) nucleic acids encoding an aforementioned peptide; (iv) administration of antisense nucleic acid and nucleic acids that are "dysfunctional" (*i.e.*, due to a heterologous insertion within the coding sequences of coding sequences to an aforementioned peptide) that are utilized to "knockout" endogenous function of an aforementioned peptide by homologous recombination (see, *e.g.*, Capecchi, 1989. Science 244: 1288-1292); or (v) modulators (*i.e.*, inhibitors, agonists and antagonists, including additional peptide mimetic of the invention or antibodies specific to a peptide of the invention) that alter the interaction between an aforementioned peptide and its binding partner.

Diseases and disorders that are characterized by decreased (relative to a subject not suffering from the disease or disorder) levels or biological activity may be treated with

Therapeutics that increase (*i.e.*, are agonists to) activity. Therapeutics that upregulate activity may be administered in a therapeutic or prophylactic manner. Therapeutics that may be utilized include, but are not limited to, an aforementioned peptide, or analogs, derivatives, fragments or homologs thereof; or an agonist that increases bioavailability.

5 Increased or decreased levels can be readily detected by quantifying peptide and/or RNA, by obtaining a patient tissue sample (*e.g.*, from biopsy tissue) and assaying it *in vitro* for RNA or peptide levels, structure and/or activity of the expressed peptides (or mRNAs of an aforementioned peptide). Methods that are well-known within the art include, but are not limited to, immunoassays (*e.g.*, by Western blot analysis, immunoprecipitation followed by
10 sodium dodecyl sulfate (SDS) polyacrylamide gel electrophoresis, immunocytochemistry, etc.) and/or hybridization assays to detect expression of mRNAs (*e.g.*, Northern assays, dot blots, in situ hybridization, etc.).

Prophylactic Methods

15 In one aspect, the invention provides a method for preventing, in a subject, a disease or condition associated with an aberrant FCTR_X expression or activity, by administering to the subject an agent that modulates FCTR_X expression or at least one FCTR_X activity. Subjects at risk for a disease that is caused or contributed to by aberrant FCTR_X expression or activity can be identified by, for example, any or a combination of diagnostic or prognostic assays as
20 described herein. Administration of a prophylactic agent can occur prior to the manifestation of symptoms characteristic of the FCTR_X aberrancy, such that a disease or disorder is prevented or, alternatively, delayed in its progression. Depending on the type of FCTR_X aberrancy, for example, an FCTR_X agonist or FCTR_X antagonist agent can be used for treating the subject. The appropriate agent can be determined based on screening assays
25 described herein. The prophylactic methods of the present invention are further discussed in the following subsections.

Therapeutic Methods

30 Another aspect of the invention pertains to methods of modulating FCTR_X expression or activity for therapeutic purposes. The modulatory method of the invention involves contacting a cell with an agent that modulates one or more of the activities of FCTR_X protein activity associated with the cell. An agent that modulates FCTR_X protein activity can be an agent as described herein, such as a nucleic acid or a protein, a naturally-occurring cognate ligand of an FCTR_X protein, a peptide, an FCTR_X peptidomimetic, or other small molecule.

In one embodiment, the agent stimulates one or more FCTR_X protein activity. Examples of such stimulatory agents include active FCTR_X protein and a nucleic acid molecule encoding FCTR_X that has been introduced into the cell. In another embodiment, the agent inhibits one or more FCTR_X protein activity. Examples of such inhibitory agents include antisense FCTR_X nucleic acid molecules and anti-FCTR_X antibodies. These modulatory methods can be performed *in vitro* (e.g., by culturing the cell with the agent) or, alternatively, *in vivo* (e.g., by administering the agent to a subject). As such, the present invention provides methods of treating an individual afflicted with a disease or disorder characterized by aberrant expression or activity of an FCTR_X protein or nucleic acid molecule. In one embodiment, the method involves administering an agent (e.g., an agent identified by a screening assay described herein), or combination of agents that modulates (e.g., upregulates or downregulates) FCTR_X expression or activity. In another embodiment, the method involves administering an FCTR_X protein or nucleic acid molecule as therapy to compensate for reduced or aberrant FCTR_X expression or activity.

Stimulation of FCTR_X activity is desirable in situations in which FCTR_X is abnormally downregulated and/or in which increased FCTR_X activity is likely to have a beneficial effect. One example of such a situation is where a subject has a disorder characterized by aberrant cell proliferation and/or differentiation (e.g., cancer). Another example of such a situation is where the subject has a gestational disease (e.g., preeclampsia).

DISEASE PATHWAYS

Determination of the Biological Effect of the Therapeutic

In various embodiments of the present invention, suitable *in vitro* or *in vivo* assays are utilized to determine the effect of a specific Therapeutic and whether its administration is indicated for treatment of the affected tissue.

In various specific embodiments, *in vitro* assays may be performed with representative cells of the type(s) involved in the patient's disorder, to determine if a given Therapeutic exerts the desired effect upon the cell type(s). Compounds for use in therapy may be tested in suitable animal model systems including, but not limited to rats, mice, chicken, cows, monkeys, rabbits, and the like, prior to testing in human subjects. Similarly, for *in vivo* testing, any of the animal model systems known in the art may be used prior to administration to human subjects.

Malignancies

An aforementioned protein is involved in the regulation of cell proliferation. Accordingly, Therapeutics of the present invention are useful in the therapeutic or prophylactic treatment of diseases or disorders that are associated with cell hyperproliferation and/or loss of control of cell proliferation (*e.g.*, cancers, malignancies and tumors). For a review of such hyperproliferation disorders, see *e.g.*, Fishman, *et al.*, 1985. MEDICINE, 2nd ed., J.B. Lippincott Co., Philadelphia, PA.

Therapeutics of the present invention may be assayed by any method known within the art for efficacy in treating or preventing malignancies and related disorders. Such assays include, but are not limited to, *in vitro* assays utilizing transformed cells or cells derived from the patient's tumor, as well as *in vivo* assays using animal models of cancer or malignancies. Potentially effective Therapeutics are those that, for example, inhibit the proliferation of tumor-derived or transformed cells in culture or cause a regression of tumors in animal models, in comparison to the controls.

In the practice of the present invention, once a malignancy or cancer has been shown to be amenable to treatment by modulating (*i.e.*, inhibiting, antagonizing or agonizing) activity, that cancer or malignancy may subsequently be treated or prevented by the administration of a Therapeutic that serves to modulate protein function.

Premalignant conditions

The Therapeutics of the present invention that are effective in the therapeutic or prophylactic treatment of cancer or malignancies may also be administered for the treatment of pre-malignant conditions and/or to prevent the progression of a pre-malignancy to a neoplastic or malignant state. Such prophylactic or therapeutic use is indicated in conditions known or suspected of preceding progression to neoplasia or cancer, in particular, where non-neoplastic cell growth consisting of hyperplasia, metaplasia or, most particularly, dysplasia has occurred. For a review of such abnormal cell growth see *e.g.*, Robbins & Angell, 1976. BASIC PATHOLOGY, 2nd ed., W.B. Saunders Co., Philadelphia, PA.

Hyperplasia is a form of controlled cell proliferation involving an increase in cell number in a tissue or organ, without significant alteration in its structure or function. For example, it has been demonstrated that endometrial hyperplasia often precedes endometrial cancer. Metaplasia is a form of controlled cell growth in which one type of mature or fully differentiated cell substitutes for another type of mature cell. Metaplasia may occur in epithelial or connective tissue cells. Dysplasia is generally considered a precursor of cancer, and is found mainly in the epithelia. Dysplasia is the most disorderly form of non-neoplastic

cell growth, and involves a loss in individual cell uniformity and in the architectural orientation of cells. Dysplasia characteristically occurs where there exists chronic irritation or inflammation, and is often found in the cervix, respiratory passages, oral cavity, and gall bladder.

5 Alternatively, or in addition to the presence of abnormal cell growth characterized as hyperplasia, metaplasia, or dysplasia, the presence of one or more characteristics of a transformed or malignant phenotype displayed either *in vivo* or *in vitro* within a cell sample derived from a patient, is indicative of the desirability of prophylactic/therapeutic administration of a Therapeutic that possesses the ability to modulate activity of An
10 aforementioned protein. Characteristics of a transformed phenotype include, but are not limited to: (i) morphological changes; (ii) looser substratum attachment; (iii) loss of cell-to-cell contact inhibition; (iv) loss of anchorage dependence; (v) protease release; (vi) increased sugar transport; (vii) decreased serum requirement; (viii) expression of fetal antigens, (ix) disappearance of the 250 kDal cell-surface protein, and the like. See *e.g.*,
15 Richards, *et al.*, 1986. MOLECULAR PATHOLOGY, W.B. Saunders Co., Philadelphia, PA.

 In a specific embodiment of the present invention, a patient that exhibits one or more of the following predisposing factors for malignancy is treated by administration of an effective amount of a Therapeutic: (i) a chromosomal translocation associated with a malignancy (*e.g.*, the Philadelphia chromosome (*bcr/abl*) for chronic myelogenous leukemia and t(14;18) for follicular lymphoma, etc.); (ii) familial polyposis or Gardner's syndrome
20 (possible forerunners of colon cancer); (iii) monoclonal gammopathy of undetermined significance (a possible precursor of multiple myeloma) and (iv) a first degree kinship with persons having a cancer or pre-cancerous disease showing a Mendelian (genetic) inheritance pattern (*e.g.*, familial polyposis of the colon, Gardner's syndrome, hereditary exostosis,
25 polyendocrine adenomatosis, Peutz-Jeghers syndrome, neurofibromatosis of Von Recklinghausen, medullary thyroid carcinoma with amyloid production and pheochromocytoma, retinoblastoma, carotid body tumor, cutaneous melanocarcinoma, intraocular melanocarcinoma, xeroderma pigmentosum, ataxia telangiectasia, Chediak-Higashi syndrome, albinism, Fanconi's aplastic anemia and Bloom's syndrome).

30 In another embodiment, a Therapeutic of the present invention is administered to a human patient to prevent the progression to breast, colon, lung, pancreatic, or uterine cancer, or melanoma or sarcoma.

Hyperproliferative and dysproliferative disorders

In one embodiment of the present invention, a Therapeutic is administered in the therapeutic or prophylactic treatment of hyperproliferative or benign dysproliferative disorders. The efficacy in treating or preventing hyperproliferative diseases or disorders of a
5 Therapeutic of the present invention may be assayed by any method known within the art. Such assays include *in vitro* cell proliferation assays, *in vitro* or *in vivo* assays using animal models of hyperproliferative diseases or disorders, or the like. Potentially effective Therapeutics may, for example, promote cell proliferation in culture or cause growth or cell proliferation in animal models in comparison to controls.

10 Specific embodiments of the present invention are directed to the treatment or prevention of cirrhosis of the liver (a condition in which scarring has overtaken normal liver regeneration processes); treatment of keloid (hypertrophic scar) formation causing disfiguring of the skin in which the scarring process interferes with normal renewal; psoriasis (a common skin condition characterized by excessive proliferation of the skin and delay in proper cell fate
15 determination); benign tumors; fibrocystic conditions and tissue hypertrophy (e.g., benign prostatic hypertrophy).

Neurodegenerative disorders

FCTR_X has been implicated in the deregulation of cellular maturation and apoptosis, which are both characteristic of neurodegenerative disease. Accordingly, Therapeutics of the
20 invention, particularly but not limited to those that modulate (or supply) activity of an aforementioned protein, may be effective in treating or preventing neurodegenerative disease. Therapeutics of the present invention that modulate the activity of an aforementioned protein involved in neurodegenerative disorders can be assayed by any method known in the art for efficacy in treating or preventing such neurodegenerative diseases and disorders. Such assays
25 include *in vitro* assays for regulated cell maturation or inhibition of apoptosis or *in vivo* assays using animal models of neurodegenerative diseases or disorders, or any of the assays described below. Potentially effective Therapeutics, for example but not by way of limitation, promote regulated cell maturation and prevent cell apoptosis in culture, or reduce neurodegeneration in animal models in comparison to controls.

30 Once a neurodegenerative disease or disorder has been shown to be amenable to treatment by modulation activity, that neurodegenerative disease or disorder can be treated or prevented by administration of a Therapeutic that modulates activity. Such diseases include all degenerative disorders involved with aging, especially osteoarthritis and neurodegenerative disorders.

Disorders related to organ transplantation

FCTR_X has been implicated in disorders related to organ transplantation, in particular but not limited to organ rejection. Therapeutics of the invention, particularly those that modulate (or supply) activity, may be effective in treating or preventing diseases or disorders related to organ transplantation. Therapeutics of the invention (particularly Therapeutics that modulate the levels or activity of an aforementioned protein) can be assayed by any method known in the art for efficacy in treating or preventing such diseases and disorders related to organ transplantation. Such assays include *in vitro* assays for using cell culture models as described below, or *in vivo* assays using animal models of diseases and disorders related to organ transplantation, see *e.g.*, below. Potentially effective Therapeutics, for example but not by way of limitation, reduce immune rejection responses in animal models in comparison to controls.

Accordingly, once diseases and disorders related to organ transplantation are shown to be amenable to treatment by modulation of activity, such diseases or disorders can be treated or prevented by administration of a Therapeutic that modulates activity.

Cardiovascular Disease

FCTR_X has been implicated in cardiovascular disorders, including in atherosclerotic plaque formation. Diseases such as cardiovascular disease, including cerebral thrombosis or hemorrhage, ischemic heart or renal disease, peripheral vascular disease, or thrombosis of other major vessel, and other diseases, including diabetes mellitus, hypertension, hypothyroidism, cholesterol ester storage disease, systemic lupus erythematosus, homocysteinemia, and familial protein or lipid processing diseases, and the like, are either directly or indirectly associated with atherosclerosis. Accordingly, Therapeutics of the invention, particularly those that modulate (or supply) activity or formation may be effective in treating or preventing atherosclerosis-associated diseases or disorders. Therapeutics of the invention (particularly Therapeutics that modulate the levels or activity) can be assayed by any method known in the art, including those described below, for efficacy in treating or preventing such diseases and disorders.

A vast array of animal and cell culture models exist for processes involved in atherosclerosis. A limited and non-exclusive list of animal models includes knockout mice for premature atherosclerosis (Kurabayashi and Yazaki, 1996, *Int. Angiol.* 15: 187-194), transgenic mouse models of atherosclerosis (Kappel *et al.*, 1994, *FASEB J.* 8: 583-592), antisense oligonucleotide treatment of animal models (Callow, 1995, *Curr. Opin. Cardiol.* 10: 569-576), transgenic rabbit models for atherosclerosis (Taylor, 1997, *Ann. N.Y. Acad.*

Sci 811: 146-152), hypercholesterolemic animal models (Rosenfeld, 1996, Diabetes Res. Clin. Pract. 30 Suppl.: 1-11), hyperlipidemic mice (Paigen *et al.*, 1994, Curr. Opin. Lipidol. 5: 258-264), and inhibition of lipoxygenase in animals (Sigal *et al.*, 1994, Ann. N.Y. Acad. Sci. 714: 211-224). In addition, *in vitro* cell models include but are not limited to monocytes
5 exposed to low density lipoprotein (Frostegard *et al.*, 1996, Atherosclerosis 121: 93-103),
cloned vascular smooth muscle cells (Suttles *et al.*, 1995, Exp. Cell Res. 218: 331-338),
endothelial cell-derived chemoattractant exposed T cells (Katz *et al.*, 1994, J. Leukoc. Biol.
55: 567-573), cultured human aortic endothelial cells (Farber *et al.*, 1992, Am. J. Physiol.
262: H1088-1085), and foam cell cultures (Libby *et al.*, 1996, Curr Opin Lipidol 7: 330-335).
10 Potentially effective Therapeutics, for example but not by way of limitation, reduce foam cell
formation in cell culture models, or reduce atherosclerotic plaque formation in
hypercholesterolemic mouse models of atherosclerosis in comparison to controls.

Accordingly, once an atherosclerosis-associated disease or disorder has been shown to
be amenable to treatment by modulation of activity or formation, that disease or disorder can
15 be treated or prevented by administration of a Therapeutic that modulates activity.

Cytokine and Cell Proliferation/Differentiation Activity

An FCTR protein of the present invention may exhibit cytokine, cell proliferation
(either inducing or inhibiting) or cell differentiation (either inducing or inhibiting) activity or
may induce production of other cytokines in certain cell populations. Many protein factors
20 discovered to date, including all known cytokines, have exhibited activity in one or more
factor dependent cell proliferation assays, and hence the assays serve as a convenient
confirmation of cytokine activity. The activity of a protein of the present invention is
evidenced by any one of a number of routine factor dependent cell proliferation assays for cell
lines including, without limitation, 32D, DA2, DA1G, T10, B9, B9/11, BaF3, MC9/G, M+
25 (preB M+), 2E8, RB5, DA1, 123, T1165, HT2, CTLL2, TF-1, Mo7e and CMK.

The activity of a protein of the invention may, among other means, be measured by the
following methods: Assays for T-cell or thymocyte proliferation include without limitation
those described in: CURRENT PROTOCOLS IN IMMUNOLOGY, Ed by Coligan *et al.*, Greene
Publishing Associates and Wiley-Interscience (Chapter 3 and Chapter 7); Takai *et al.*, *J*
30 *Immunol* 137:3494-3500, 1986; Bertagnoli *et al.*, *J Immunol* 145:1706-1712, 1990;
Bertagnoli *et al.*, *Cell Immunol* 133:327-341, 1991; Bertagnoli, *et al.*, *J Immunol*
149:3778-3783, 1992; Bowman *et al.*, *J Immunol* 152:1756-1761, 1994.

Assays for cytokine production and/or proliferation of spleen cells, lymph node cells or
thymocytes include, without limitation, those described by Kruisbeek and Shevach, In:

CURRENT PROTOCOLS IN IMMUNOLOGY. Coligan *et al.*, eds., Vol 1, pp. 3.12.1-14, John Wiley and Sons, Toronto 1994; and by Schreiber, In: CURRENT PROTOCOLS IN IMMUNOLOGY. Coligan eds., Vol 1, pp. 6.8.1-8, John Wiley and Sons, Toronto 1994.

Assays for proliferation and differentiation of hematopoietic and lymphopoietic cells include, without limitation, those described by Bottomly *et al.*, In: CURRENT PROTOCOLS IN IMMUNOLOGY. Coligan *et al.*, eds., Vol 1, pp. 6.3.1-6.3.12, John Wiley and Sons, Toronto 1991; deVries *et al.*, *J Exp Med* 173:1205-1211, 1991; Moreau *et al.*, *Nature* 336:690-692, 1988; Greenberger *et al.*, *Proc Natl Acad Sci U.S.A.* 80:2931-2938, 1983; Nordan, In: CURRENT PROTOCOLS IN IMMUNOLOGY. Coligan *et al.*, eds., Vol 1, pp. 6.6.1-5, John Wiley and Sons, Toronto 1991; Smith *et al.*, *Proc Natl Acad Sci U.S.A.* 83:1857-1861, 1986; Measurement of human Interleukin 11-Bennett, *et al.* In: CURRENT PROTOCOLS IN IMMUNOLOGY. Coligan *et al.*, eds., Vol 1, pp. 6.15.1 John Wiley and Sons, Toronto 1991; Ciarletta, *et al.*, In: CURRENT PROTOCOLS IN IMMUNOLOGY. Coligan *et al.*, eds., Vol 1, pp. 6.13.1, John Wiley and Sons, Toronto 1991.

Assays for T-cell clone responses to antigens (which will identify, among others, proteins that affect APC-T cell interactions as well as direct T-cell effects by measuring proliferation and cytokine production) include, without limitation, those described In: CURRENT PROTOCOLS IN IMMUNOLOGY. Coligan *et al.*, eds., Greene Publishing Associates and Wiley-Interscience (Chapter 3, Chapter 6, Chapter 7); Weinberger *et al.*, *Proc Natl Acad Sci USA* 77:6091-6095, 1980; Weinberger *et al.*, *Eur J Immun* 11:405-411, 1981; Takai *et al.*, *J Immunol* 137:3494-3500, 1986; Takai *et al.*, *J Immunol* 140:508-512, 1988.

Immune Stimulating or Suppressing Activity

An FCTR protein of the present invention may also exhibit immune stimulating or immune suppressing activity, including without limitation the activities for which assays are described herein. A protein are useful in the treatment of various immune deficiencies and disorders (including severe combined immunodeficiency (SCID)), *e.g.*, in regulating (up or down) growth and proliferation of T and/or B lymphocytes, as well as effecting the cytolytic activity of NK cells and other cell populations. These immune deficiencies may be genetic or be caused by viral (*e.g.*, HIV) as well as bacterial or fungal infections, or may result from autoimmune disorders. More specifically, infectious diseases caused by viral, bacterial, fungal or other infection may be treatable using a protein of the present invention, including infections by HIV, hepatitis viruses, herpesviruses, mycobacteria, Leishmania species, malaria species, and various fungal infections such as candidiasis. Of course, in this regard, a protein

of the present invention may also be useful where a boost to the immune system generally may be desirable, *i.e.*, in the treatment of cancer.

Autoimmune disorders which may be treated using a protein of the present invention include, for example, connective tissue disease, multiple sclerosis, systemic lupus
5 erythematosus, rheumatoid arthritis, autoimmune pulmonary inflammation, Guillain-Barre syndrome, autoimmune thyroiditis, insulin dependent diabetes mellitus, myasthenia gravis, graft-versus-host disease and autoimmune inflammatory eye disease. Such a protein of the present invention may also to be useful in the treatment of allergic reactions and conditions, such as asthma (particularly allergic asthma) or other respiratory problems. Other conditions,
10 in which immune suppression is desired (including, for example, organ transplantation), may also be treatable using a protein of the present invention.

Using the proteins of the invention it may also be possible to immune responses, in a number of ways. Down regulation may be in the form of inhibiting or blocking an immune response already in progress or may involve preventing the induction of an immune response.
15 The functions of activated T cells may be inhibited by suppressing T cell responses or by inducing specific tolerance in T cells, or both. Immunosuppression of T cell responses is generally an active, non-antigen-specific, process which requires continuous exposure of the T cells to the suppressive agent. Tolerance, which involves inducing non-responsiveness or energy in T cells, is distinguishable from immunosuppression in that it is generally
20 antigen-specific and persists after exposure to the tolerizing agent has ceased. Operationally, tolerance can be demonstrated by the lack of a T cell response upon re-exposure to specific antigen in the absence of the tolerizing agent.

Down regulating or preventing one or more antigen functions (including without limitation B lymphocyte antigen functions (such as, for example, B7), *e.g.*, preventing high
25 level lymphokine synthesis by activated T cells, are useful in situations of tissue, skin and organ transplantation and in graft-versus-host disease (GVHD). For example, blockage of T cell function should result in reduced tissue destruction in tissue transplantation. Typically, in tissue transplants, rejection of the transplant is initiated through its recognition as foreign by T cells, followed by an immune reaction that destroys the transplant. The administration of a
30 molecule which inhibits or blocks interaction of a B7 lymphocyte antigen with its natural ligand(s) on immune cells (such as a soluble, monomeric form of a peptide having B7-2 activity alone or in conjunction with a monomeric form of a peptide having an activity of another B lymphocyte antigen (*e.g.*, B7-1, B7-3) or blocking antibody), prior to transplantation can lead to the binding of the molecule to the natural ligand(s) on the immune cells without

transmitting the corresponding costimulatory signal. Blocking B lymphocyte antigen function in this matter prevents cytokine synthesis by immune cells, such as T cells, and thus acts as an immunosuppressant. Moreover, the lack of costimulation may also be sufficient to energize the T cells, thereby inducing tolerance in a subject. Induction of long-term tolerance by B lymphocyte antigen-blocking reagents may avoid the necessity of repeated administration of these blocking reagents. To achieve sufficient immunosuppression or tolerance in a subject, it may also be necessary to block the function of B lymphocyte antigens.

The efficacy of particular blocking reagents in preventing organ transplant rejection or GVHD can be assessed using animal models that are predictive of efficacy in humans. Examples of appropriate systems which can be used include allogeneic cardiac grafts in rats and xenogeneic pancreatic islet cell grafts in mice, both of which have been used to examine the immunosuppressive effects of CTLA4Ig fusion proteins *in vivo* as described in Lenschow *et al.*, Science 257:789-792 (1992) and Turka *et al.*, Proc Natl Acad Sci USA, 89:11102-11105 (1992). In addition, murine models of GVHD (see Paul ed., FUNDAMENTAL IMMUNOLOGY, Raven Press, New York, 1989, pp. 846-847) can be used to determine the effect of blocking B lymphocyte antigen function *in vivo* on the development of that disease.

Blocking antigen function may also be therapeutically useful for treating autoimmune diseases. Many autoimmune disorders are the result of inappropriate activation of T cells that are reactive against self tissue and which promote the production of cytokines and auto-antibodies involved in the pathology of the diseases. Preventing the activation of autoreactive T cells may reduce or eliminate disease symptoms. Administration of reagents which block costimulation of T cells by disrupting receptor:ligand interactions of B lymphocyte antigens can be used to inhibit T cell activation and prevent production of auto-antibodies or T cell-derived cytokines which may be involved in the disease process. Additionally, blocking reagents may induce antigen-specific tolerance of autoreactive T cells which could lead to long-term relief from the disease. The efficacy of blocking reagents in preventing or alleviating autoimmune disorders can be determined using a number of well-characterized animal models of human autoimmune diseases. Examples include murine experimental autoimmune encephalitis, systemic lupus erythematosus in MRL/lpr/lpr mice or NZB hybrid mice, murine autoimmune collagen arthritis, diabetes mellitus in NOD mice and BB rats, and murine experimental myasthenia gravis (see Paul ed., FUNDAMENTAL IMMUNOLOGY, Raven Press, New York, 1989, pp. 840-856).

Upregulation of an antigen function (preferably a B lymphocyte antigen function), as a means of up regulating immune responses, may also be useful in therapy. Upregulation of

immune responses may be in the form of enhancing an existing immune response or eliciting an initial immune response. For example, enhancing an immune response through stimulating B lymphocyte antigen function are useful in cases of viral infection. In addition, systemic viral diseases such as influenza, the common cold, and encephalitis might be alleviated by the administration of stimulatory forms of B lymphocyte antigens systemically.

Alternatively, anti-viral immune responses may be enhanced in an infected patient by removing T cells from the patient, costimulating the T cells *in vitro* with viral antigen-pulsed APCs either expressing a peptide of the present invention or together with a stimulatory form of a soluble peptide of the present invention and reintroducing the *in vitro* activated T cells into the patient. Another method of enhancing anti-viral immune responses would be to isolate infected cells from a patient, transfect them with a nucleic acid encoding a protein of the present invention as described herein such that the cells express all or a portion of the protein on their surface, and reintroduce the transfected cells into the patient. The infected cells would now be capable of delivering a costimulatory signal to, and thereby activate, T cells *in vivo*.

In another application, up regulation or enhancement of antigen function (preferably B lymphocyte antigen function) are useful in the induction of tumor immunity. Tumor cells (*e.g.*, sarcoma, melanoma, lymphoma, leukemia, neuroblastoma, carcinoma) transfected with a nucleic acid encoding at least one peptide of the present invention can be administered to a subject to overcome tumor-specific tolerance in the subject. If desired, the tumor cell can be transfected to express a combination of peptides. For example, tumor cells obtained from a patient can be transfected *ex vivo* with an expression vector directing the expression of a peptide having B7-2-like activity alone, or in conjunction with a peptide having B7-1-like activity and/or B7-3-like activity. The transfected tumor cells are returned to the patient to result in expression of the peptides on the surface of the transfected cell. Alternatively, gene therapy techniques can be used to target a tumor cell for transfection *in vivo*.

The presence of the peptide of the present invention having the activity of a B lymphocyte antigen(s) on the surface of the tumor cell provides the necessary costimulation signal to T cells to induce a T cell mediated immune response against the transfected tumor cells. In addition, tumor cells which lack MHC class I or MHC class II molecules, or which fail to reexpress sufficient amounts of MHC class I or MHC class II molecules, can be transfected with nucleic acid encoding all or a portion of (*e.g.*, a cytoplasmic-domain truncated portion) of an MHC class I a chain protein and b 2 microglobulin protein or an MHC class II a chain protein and an MHC class II b chain protein to thereby express MHC class I or MHC

class II proteins on the cell surface. Expression of the appropriate class I or class II MHC in conjunction with a peptide having the activity of a B lymphocyte antigen (e.g., B7-1, B7-2, B7-3) induces a T cell mediated immune response against the transfected tumor cell.

Optionally, a gene encoding an antisense construct which blocks expression of an MHC class II associated protein, such as the invariant chain, can also be cotransfected with a DNA encoding a peptide having the activity of a B lymphocyte antigen to promote presentation of tumor associated antigens and induce tumor specific immunity. Thus, the induction of a T cell mediated immune response in a human subject may be sufficient to overcome tumor-specific tolerance in the subject.

The activity of a protein of the invention may, among other means, be measured by the following methods: Suitable assays for thymocyte or splenocyte cytotoxicity include, without limitation, those described in: CURRENT PROTOCOLS IN IMMUNOLOGY. Coligan *et al.*, eds. Greene Publishing Associates and Wiley-Interscience (Chapter 3, Chapter 7); Herrmann *et al.*, *Proc Natl Acad Sci USA* 78:2488-2492, 1981; Herrmann *et al.*, *J Immunol* 128:1968-1974, 1982; Handa *et al.*, *J Immunol* 135:1564-1572, 1985; Takai *et al.*, *J Immunol* 137:3494-3500, 1986; Takai *et al.*, *J Immunol* 140:508-512, 1988; Herrmann *et al.*, *Proc Natl Acad Sci USA* 78:2488-2492, 1981; Herrmann *et al.*, *J Immunol* 128:1968-1974, 1982; Handa *et al.*, *J Immunol* 135:1564-1572, 1985; Takai *et al.*, *J Immunol* 137:3494-3500, 1986; Bowman *et al.*, *J Virology* 61:1992-1998; Takai *et al.*, *J Immunol* 140:508-512, 1988; Bertagnolli *et al.*, *Cell Immunol* 133:327-341, 1991; Brown *et al.*, *J Immunol* 153:3079-3092, 1994.

Assays for T-cell-dependent immunoglobulin responses and isotype switching (which will identify, among others, proteins that modulate T-cell dependent antibody responses and that affect Th1/Th2 profiles) include, without limitation, those described in: Maliszewski, *J Immunol* 144:3028-3033, 1990; and Mond and Brunswick In: CURRENT PROTOCOLS IN IMMUNOLOGY. Coligan *et al.*, eds., Vol 1, pp. 3.8.1-3.8.16, John Wiley and Sons, Toronto 1994.

Mixed lymphocyte reaction (MLR) assays (which will identify, among others, proteins that generate predominantly Th1 and CTL responses) include, without limitation, those described in: CURRENT PROTOCOLS IN IMMUNOLOGY. Coligan *et al.*, eds. Greene Publishing Associates and Wiley-Interscience (Chapter 3, Chapter 7); Takai *et al.*, *J Immunol* 137:3494-3500, 1986; Takai *et al.*, *J Immunol* 140:508-512, 1988; Bertagnolli *et al.*, *J Immunol* 149:3778-3783, 1992.

Dendritic cell-dependent assays (which will identify, among others, proteins expressed by dendritic cells that activate naive T-cells) include, without limitation, those described in:

Guery *et al.*, *J Immunol* 134:536-544, 1995; Inaba *et al.*, *J Exp Med* 173:549-559, 1991; Macatonia *et al.*, *J Immunol* 154:5071-5079, 1995; Porgador *et al.*, *J Exp Med* 182:255-260, 1995; Nair *et al.*, *J Virol* 67:4062-4069, 1993; Huang *et al.*, *Science* 264:961-965, 1994; Macatonia *et al.*, *J Exp Med* 169:1255-1264, 1989; Bhardwaj *et al.*, *J Clin Investig* 5 94:797-807, 1994; and Inaba *et al.*, *J Exp Med* 172:631-640, 1990.

Assays for lymphocyte survival/apoptosis (which will identify, among others, proteins that prevent apoptosis after superantigen induction and proteins that regulate lymphocyte homeostasis) include, without limitation, those described in: Darzynkiewicz *et al.*, *Cytometry* 13:795-808, 1992; Gorczyca *et al.*, *Leukemia* 7:659-670, 1993; Gorczyca *et al.*, *Cancer Res* 10 53:1945-1951, 1993; Itoh *et al.*, *Cell* 66:233-243, 1991; Zacharchuk, *J Immunol* 145:4037-4045, 1990; Zamai *et al.*, *Cytometry* 14:891-897, 1993; Gorczyca *et al.*, *Internat J Oncol* 1:639-648, 1992.

Assays for proteins that influence early steps of T-cell commitment and development include, without limitation, those described in: Antica *et al.*, *Blood* 84:111-117, 1994; Fine *et al.*, *Cell Immunol* 155: 111-122, 1994; Galy *et al.*, *Blood* 85:2770-2778, 1995; Toki *et al.*, 15 *Proc Nat Acad Sci USA* 88:7548-7551, 1991.

Hematopoiesis Regulating Activity

An FCTR protein of the present invention are useful in regulation of hematopoiesis and, consequently, in the treatment of myeloid or lymphoid cell deficiencies. Even marginal biological activity in support of colony forming cells or of factor-dependent cell lines indicates involvement in regulating hematopoiesis, *e.g.* in supporting the growth and proliferation of erythroid progenitor cells alone or in combination with other cytokines, thereby indicating utility, for example, in treating various anemias or for use in conjunction with 20 irradiation/chemotherapy to stimulate the production of erythroid precursors and/or erythroid cells; in supporting the growth and proliferation of myeloid cells such as granulocytes and monocytes/macrophages (*i.e.*, traditional CSF activity) useful, for example, in conjunction with chemotherapy to prevent or treat consequent myelo-suppression; in supporting the growth and proliferation of megakaryocytes and consequently of platelets thereby allowing prevention 25 or treatment of various platelet disorders such as thrombocytopenia, and generally for use in place of or complimentary to platelet transfusions; and/or in supporting the growth and proliferation of hematopoietic stem cells which are capable of maturing to any and all of the above-mentioned hematopoietic cells and therefore find therapeutic utility in various stem cell disorders (such as those usually treated with transplantation, including, without limitation, 30

aplastic anemia and paroxysmal nocturnal hemoglobinuria), as well as in repopulating the stem cell compartment post irradiation/chemotherapy, either in-vivo or ex-vivo (*i.e.*, in conjunction with bone marrow transplantation or with peripheral progenitor cell transplantation (homologous or heterologous)) as normal cells or genetically manipulated for gene therapy.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Suitable assays for proliferation and differentiation of various hematopoietic lines are cited above.

Assays for embryonic stem cell differentiation (which will identify, among others, proteins that influence embryonic differentiation hematopoiesis) include, without limitation, those described in: Johansson *et al.* Cellular Biology 15:141-151, 1995; Keller *et al.*, Molecular and Cellular Biology 13:473-486, 1993; McClanahan *et al.*, Blood 81:2903-2915, 1993.

Assays for stem cell survival and differentiation (which will identify, among others, proteins that regulate lympho-hematopoiesis) include, without limitation, those described in: Methylcellulose colony forming assays, Freshney, M. G. In CULTURE OF HEMATOPOIETIC CELLS. Freshney, *et al.* eds. Vol pp. 265-268, Wiley-Liss, Inc., New York, N.Y. 1994; Hirayama *et al.*, *Proc Natl Acad Sci USA* 89:5907-5911, 1992; McNiece and Briddeli. In CULTURE OF HEMATOPOIETIC CELLS. Freshney, *et al.* eds. Vol pp. 23-39, Wiley-Liss, Inc., New York, N.Y. 1994; Neben *et al.*, *Exp Hematol* 22:353-359, 1994; Ploemacher In CULTURE OF HEMATOPOIETIC CELLS. Freshney, *et al.* eds. Vol pp. 1-21, Wiley-Liss, Inc., New York, N.Y. 1994; Spoonceret *et al.*, In CULTURE OF HEMATOPOIETIC CELLS. Freshney, *et al.* eds. Vol pp. 163-179, Wiley-Liss, Inc., New York, N.Y. 1994; Sutherland, In CULTURE OF HEMATOPOIETIC CELLS. Freshney, *et al.* eds. Vol pp. 139-162, Wiley-Liss, Inc., New York, N.Y. 1994.

Tissue Growth Activity

An FCTR_X protein of the present invention also may have utility in compositions used for bone, cartilage, tendon, ligament and/or nerve tissue growth or regeneration, as well as for wound healing and tissue repair and replacement, and in the treatment of burns, incisions and ulcers.

A protein of the present invention, which induces cartilage and/or bone growth in circumstances where bone is not normally formed, has application in the healing of bone

fractures and cartilage damage or defects in humans and other animals. Such a preparation employing a protein of the invention may have prophylactic use in closed as well as open fracture reduction and also in the improved fixation of artificial joints. De novo bone formation induced by an osteogenic agent contributes to the repair of congenital, trauma induced, or oncologic resection induced craniofacial defects, and also is useful in cosmetic plastic surgery.

A protein of this invention may also be used in the treatment of periodontal disease, and in other tooth repair processes. Such agents may provide an environment to attract bone-forming cells, stimulate growth of bone-forming cells or induce differentiation of progenitors of bone-forming cells. A protein of the invention may also be useful in the treatment of osteoporosis or osteoarthritis, such as through stimulation of bone and/or cartilage repair or by blocking inflammation or processes of tissue destruction (collagenase activity, osteoclast activity, etc.) mediated by inflammatory processes.

Another category of tissue regeneration activity that may be attributable to the protein of the present invention is tendon/ligament formation. A protein of the present invention, which induces tendon/ligament-like tissue or other tissue formation in circumstances where such tissue is not normally formed, has application in the healing of tendon or ligament tears, deformities and other tendon or ligament defects in humans and other animals. Such a preparation employing a tendon/ligament-like tissue inducing protein may have prophylactic use in preventing damage to tendon or ligament tissue, as well as use in the improved fixation of tendon or ligament to bone or other tissues, and in repairing defects to tendon or ligament tissue. De novo tendon/ligament-like tissue formation induced by a composition of the present invention contributes to the repair of congenital, trauma induced, or other tendon or ligament defects of other origin, and is also useful in cosmetic plastic surgery for attachment or repair of tendons or ligaments. The compositions of the present invention may provide an environment to attract tendon- or ligament-forming cells, stimulate growth of tendon- or ligament-forming cells, induce differentiation of progenitors of tendon- or ligament-forming cells, or induce growth of tendon/ligament cells or progenitors *ex vivo* for return *in vivo* to effect tissue repair. The compositions of the invention may also be useful in the treatment of tendonitis, carpal tunnel syndrome and other tendon or ligament defects. The compositions may also include an appropriate matrix and/or sequestering agent as a carrier as is well known in the art.

The protein of the present invention may also be useful for proliferation of neural cells and for regeneration of nerve and brain tissue, *i.e.* for the treatment of central and peripheral nervous system diseases and neuropathies, as well as mechanical and traumatic disorders,

which involve degeneration, death or trauma to neural cells or nerve tissue. More specifically, a protein may be used in the treatment of diseases of the peripheral nervous system, such as peripheral nerve injuries, peripheral neuropathy and localized neuropathies, and central nervous system diseases, such as Alzheimer's, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, and Shy-Drager syndrome. Further conditions which may be treated in accordance with the present invention include mechanical and traumatic disorders, such as spinal cord disorders, head trauma and cerebrovascular diseases such as stroke. Peripheral neuropathies resulting from chemotherapy or other medical therapies may also be treatable using a protein of the invention.

10 Proteins of the invention may also be useful to promote better or faster closure of non-healing wounds, including without limitation pressure ulcers, ulcers associated with vascular insufficiency, surgical and traumatic wounds, and the like.

It is expected that a protein of the present invention may also exhibit activity for generation or regeneration of other tissues, such as organs (including, for example, pancreas, liver, intestine, kidney, skin, endothelium), muscle (smooth, skeletal or cardiac) and vascular (including vascular endothelium) tissue, or for promoting the growth of cells comprising such tissues. Part of the desired effects may be by inhibition or modulation of fibrotic scarring to allow normal tissue to regenerate. A protein of the invention may also exhibit angiogenic activity.

20 A protein of the present invention may also be useful for gut protection or regeneration and treatment of lung or liver fibrosis, reperfusion injury in various tissues, and conditions resulting from systemic cytokine damage.

A protein of the present invention may also be useful for promoting or inhibiting differentiation of tissues described above from precursor tissues or cells; or for inhibiting the growth of tissues described above.

25 The activity of a protein of the invention may, among other means, be measured by the following methods:

Assays for tissue generation activity include, without limitation, those described in: International Patent Publication No. WO95/16035 (bone, cartilage, tendon); International Patent Publication No. WO95/05846 (nerve, neuronal); International Patent Publication No. WO91/07491 (skin, endothelium).

30 Assays for wound healing activity include, without limitation, those described in: Winter, *Epidermal Wound Healing*, pp. 71-112 (Maibach, H I and Rovee, D T, eds.), Year

Activin/Inhibin Activity

5 An FCTR protein of the present invention may also exhibit activin- or inhibin-related activities. Inhibins are characterized by their ability to inhibit the release of follicle stimulating hormone (FSH), while activins are characterized by their ability to stimulate the release of follicle stimulating hormone (FSH). Thus, a protein of the present invention, alone or in heterodimers with a member of the inhibin family, are useful as a contraceptive
10 based on the ability of inhibins to decrease fertility in female mammals and decrease spermatogenesis in male mammals. Administration of sufficient amounts of other inhibins can induce infertility in these mammals. Alternatively, the protein of the invention, as a homodimer or as a heterodimer with other protein subunits of the inhibin-b group, are useful as a fertility inducing therapeutic, based upon the ability of activin molecules in stimulating
15 FSH release from cells of the anterior pituitary. See, for example, U.S. Pat. No. 4,798,885. A protein of the invention may also be useful for advancement of the onset of fertility in sexually immature mammals, so as to increase the lifetime reproductive performance of domestic animals such as cows, sheep and pigs.

 The activity of a protein of the invention may, among other means, be measured by the
20 following methods:

 Assays for activin/inhibin activity include, without limitation, those described in: Vale *et al.*, Endocrinology 91:562-572, 1972; Ling *et al.*, Nature 321:779-782, 1986; Vale *et al.*, Nature 321:776-779, 1986; Mason *et al.*, Nature 318:659-663, 1985; Forage *et al.*, *Proc Natl Acad Sci USA* 83:3091-3095, 1986.

25

Chemotactic/Chemokinetic Activity

 A protein of the present invention may have chemotactic or chemokinetic activity (*e.g.*, act as a chemokine) for mammalian cells, including, for example, monocytes, fibroblasts, neutrophils, T-cells, mast cells, eosinophils, epithelial and/or endothelial cells. Chemotactic
30 and chemokinetic proteins can be used to mobilize or attract a desired cell population to a desired site of action. Chemotactic or chemokinetic proteins provide particular advantages in treatment of wounds and other trauma to tissues, as well as in treatment of localized infections. For example, attraction of lymphocytes, monocytes or neutrophils to tumors or sites of infection may result in improved immune responses against the tumor or infecting agent.

A protein or peptide has chemotactic activity for a particular cell population if it can stimulate, directly or indirectly, the directed orientation or movement of such cell population. Preferably, the protein or peptide has the ability to directly stimulate directed movement of cells. Whether a particular protein has chemotactic activity for a population of cells can be readily determined by employing such protein or peptide in any known assay for cell chemotaxis.

The activity of a protein of the invention may, among other means, be measured by following methods:

Assays for chemotactic activity (which will identify proteins that induce or prevent chemotaxis) consist of assays that measure the ability of a protein to induce the migration of cells across a membrane as well as the ability of a protein to induce the adhesion of one cell population to another cell population. Suitable assays for movement and adhesion include, without limitation, those described in: CURRENT PROTOCOLS IN IMMUNOLOGY, Coligan *et al.*, eds. (Chapter 6.12, Measurement of alpha and beta Chemokines 6.12.1-6.12.28); Taub *et al.* *J Clin Invest* 95:1370-1376, 1995; Lind *et al.* *APMIS* 103:140-146, 1995; Muller *et al.* *J Immunol* 25: 1744-1748; Gruber *et al.* *J Immunol* 152:5860-5867, 1994; Johnston *et al.* *J Immunol* 153: 1762-1768, 1994.

Hemostatic and Thrombolytic Activity

A protein of the invention may also exhibit hemostatic or thrombolytic activity. As a result, such a protein is expected to be useful in treatment of various coagulation disorders (including hereditary disorders, such as hemophilias) or to enhance coagulation and other hemostatic events in treating wounds resulting from trauma, surgery or other causes. A protein of the invention may also be useful for dissolving or inhibiting formation of thromboses and for treatment and prevention of conditions resulting therefrom (such as, for example, infarction of cardiac and central nervous system vessels (*e.g.*, stroke).

The activity of a protein of the invention may, among other means, be measured by the following methods:

Assay for hemostatic and thrombolytic activity include, without limitation, those described in: Linet *et al.*, *J. Clin. Pharmacol.* 26:131-140, 1986; Burdick *et al.*, *Thrombosis Res.* 45:413-419, 1987; Humphrey *et al.*, *Fibrinolysis* 5:71-79 (1991); Schaub, *Prostaglandins* 35:467-474, 1988.

Receptor/Ligand Activity

A protein of the present invention may also demonstrate activity as receptors, receptor ligands or inhibitors or agonists of receptor/ligand interactions. Examples of such receptors and ligands include, without limitation, cytokine receptors and their ligands, receptor kinases and their ligands, receptor phosphatases and their ligands, receptors involved in cell—cell interactions and their ligands (including without limitation, cellular adhesion molecules (such as selectins, integrins and their ligands) and receptor/ligand pairs involved in antigen presentation, antigen recognition and development of cellular and humoral immune responses). Receptors and ligands are also useful for screening of potential peptide or small molecule inhibitors of the relevant receptor/ligand interaction. A protein of the present invention (including, without limitation, fragments of receptors and ligands) may themselves be useful as inhibitors of receptor/ligand interactions.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Suitable assays for receptor-ligand activity include without limitation those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A. M. Kruisbeek, D. H. Margulies, E. M. Shevach, W. Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 7.28, Measurement of Cellular Adhesion under static conditions 7.28.1-7.28.22), Takai *et al.*, *Proc Natl Acad Sci USA* 84:6864-6868, 1987; Bierer *et al.*, *J. Exp. Med.* 168:1145-1156, 1988; Rosenstein *et al.*, *J. Exp. Med.* 169:149-160 1989; Stoltenborg *et al.*, *J Immunol Methods* 175:59-68, 1994; Stitt *et al.*, *Cell* 80:661-670, 1995.

Anti-Inflammatory Activity

Proteins of the present invention may also exhibit anti-inflammatory activity. The anti-inflammatory activity may be achieved by providing a stimulus to cells involved in the inflammatory response, by inhibiting or promoting cell—cell interactions (such as, for example, cell adhesion), by inhibiting or promoting chemotaxis of cells involved in the inflammatory process, inhibiting or promoting cell extravasation, or by stimulating or suppressing production of other factors which more directly inhibit or promote an inflammatory response. Proteins exhibiting such activities can be used to treat inflammatory conditions including chronic or acute conditions), including without limitation inflammation associated with infection (such as septic shock, sepsis or systemic inflammatory response syndrome (SIRS)), ischemia-reperfusion injury, endotoxin lethality, arthritis,

complement-mediated hyperacute rejection, nephritis, cytokine or chemokine-induced lung injury, inflammatory bowel disease, Crohn's disease or resulting from over production of cytokines such as TNF or IL-1. Proteins of the invention may also be useful to treat anaphylaxis and hypersensitivity to an antigenic substance or material.

5

Tumor Inhibition Activity

In addition to the activities described above for immunological treatment or prevention of tumors, a protein of the invention may exhibit other anti-tumor activities. A protein may inhibit tumor growth directly or indirectly (such as, for example, via ADCC). A protein may exhibit its tumor inhibitory activity by acting on tumor tissue or tumor precursor tissue, by inhibiting formation of tissues necessary to support tumor growth (such as, for example, by inhibiting angiogenesis), by causing production of other factors, agents or cell types which inhibit tumor growth, or by suppressing, eliminating or inhibiting factors, agents or cell types which promote tumor growth.

15

Other Activities

A protein of the invention may also exhibit one or more of the following additional activities or effects: inhibiting the growth, infection or function of, or killing, infectious agents, including, without limitation, bacteria, viruses, fungi and other parasites; effecting (suppressing or enhancing) bodily characteristics, including, without limitation, height, weight, hair color, eye color, skin, fat to lean ratio or other tissue pigmentation, or organ or body part size or shape (such as, for example, breast augmentation or diminution, change in bone form or shape); effecting biorhythms or circadian cycles or rhythms; effecting the fertility of male or female subjects; effecting the metabolism, catabolism, anabolism, processing, utilization, storage or elimination of dietary fat, lipid, protein, carbohydrate, vitamins, minerals, cofactors or other nutritional factors or component(s); effecting behavioral characteristics, including, without limitation, appetite, libido, stress, cognition (including cognitive disorders), depression (including depressive disorders) and violent behaviors; providing analgesic effects or other pain reducing effects; promoting differentiation and growth of embryonic stem cells in lineages other than hematopoietic lineages; hormonal or endocrine activity; in the case of enzymes, correcting deficiencies of the enzyme and treating deficiency-related diseases; treatment of hyperproliferative disorders (such as, for example, psoriasis); immunoglobulin-like activity (such as, for example, the ability to bind antigens or complement); and the ability to act as an antigen in a vaccine composition to raise an immune

response against such protein or another material or entity which is cross-reactive with such protein.

The present invention is not to be limited in scope by the specific embodiments described herein. Indeed, various modifications of the invention in addition to those described
5 herein will become apparent to those skilled in the art from the foregoing description and accompanying figures. Such modifications are intended to fall within the scope of the appended claims.

This invention is further illustrated by the following examples, which should not be construed as limiting. The contents of all references, patents and published patent applications
10 cited throughout this application are hereby incorporated by reference.

Example 1. Identification of murine clone 7971c.7-r0s0-212.2-EXT (SEQ ID NO:2).

Differential gene expression was examined in mammary tumors that spontaneously
15 arise between 1 and 5 months of age in Wnt-1 transgenic mice. Expression in mammary tumors was compared to the gene expression pattern found in the normal mammary gland of wild type age matched mice. The Wnt-1 transgenic mice were obtained from Genentech, Inc. (So. San Francisco, CA). Quantitative Expression Analysis (QEA) was performed on mRNA isolated from the tumor stage of progression in these mice (8 months). Sample preparation
20 and QEA analysis are described fully in U. S. Patent No. 5,871,697 and in Shimkets et al., Nature Biotechnology 17:198-803 (1999)), incorporated herein by reference in their entireties.

Traces for two animals are shown in Fig. 6 for each of the two sets comparing transgenic mice to wild type mice. A series of transcripts overexpressed in the transgenic mice, including one at position 212.2 (vertical marker) was found. This band was designated
25 as 7971c.7 (SEQ ID NO:1).

The sequence of murine clone 7971c.7 was extended by assembly with other murine nucleic acid fragments. The resulting assembly provided the sequence of clone 7971c.7-r0s0-212.2-EXT (SEQ ID NO:2).

30 Example 2. Identification of clone 65677221-3-frag (SEQ ID NO:5), a novel human S100 cytokine

Murine clone 7971c.7 was found to have high similarity to the human EST AA315020 (SEQ ID NO:4). This sequence includes two unspecified bases designated "N" in SEQ ID NO:4. EST AA315020 was assembled with human sequences present in CuraGen
35 Corporation's sequence database obtained by its SeqCalling technology. SeqCalling is a differential expression and sequencing procedure that normalizes mRNA species in a sample,

and is disclosed in U. S. Ser. No. 09/417,386, filed Oct. 13, 1999, incorporated herein by reference in its entirety.

This process was used to assemble the sequence of SEQ ID NO:5 encompassing the sequence of SEQ ID NO:4, including providing base calls for the two unidentified bases in
5 SEQ ID NO:4. Three separate SeqCalling fragments were identified in this search, two arising from embryonic tumor cell lines (CuraGen Nos. 61097801 and 61124196), and one from endothelial cells (primary dermal endothelial cell from Cell Application, San Diego, CA; CuraGen No. 60920173) that were treated with the cytokines IL-1beta (2ng/ml) and TNF-alpha (5ng/ml) for 16hrs in order to induce leukocyte adherence. The alignment of the
10 resulting assembly with the original EST AA315020 (SEQ ID NO:4) to provide the consensus sequence identified as SEQ ID NO:5 is shown in Fig. 7. This consensus was obtained using the CAP alignment program (Huang X, Genomics 1992 Sep;14(1):18-25).

Example 3. Quantitative expression analysis of human clone 65677221-3-frag (SEQ ID NO:5) in various cells and tissues
15

The quantitative expression of various clones was assessed in about 41 normal and about 55 tumor samples by real time quantitative PCR (TAQMAN[®]) performed on a Perkin-Elmer Biosystems ABI PRISM[®] 7700 Sequence Detection System.

First, 96 RNA samples were normalized to β -actin and GAPDH. RNA (~50 ng total or
20 ~1 ng polyA+) was converted to cDNA using the TAQMAN[®] Reverse Transcription Reagents Kit (PE Biosystems, Foster City, CA; Catalog No. N808-0234) and random hexamers according to the manufacturer's protocol. Reactions were performed in 20 ul and incubated for 30 min. at 48[°]C. cDNA (5 ul) was then transferred to a separate plate for the TAQMAN[®] reaction using β -actin and GAPDH TAQMAN[®] Assay Reagents (PE Biosystems; Catalog
25 Nos. 4310881E and 4310884E, respectively) and TAQMAN[®] universal PCR Master Mix (PE Biosystems; Catalog No. 4304447) according to the manufacturer's protocol. Reactions were performed in 25 ul using the following parameters: 2 min. at 50[°]C; 10 min. at 95[°]C; 15 sec. at 95[°]C/1 min. at 60[°]C (40 cycles). Results were recorded as CT values (cycle at which a given sample crosses a threshold level of fluorescence) using a log scale, with the difference in RNA
30 concentration between a given sample and the sample with the lowest CT value being represented as 2 to the power of delta CT. The percent relative expression is then obtained by taking the reciprocal of this RNA difference and multiplying by 100. The average CT values obtained for β -actin and GAPDH were used to normalize RNA samples. The RNA sample

generating the highest CT value required no further diluting, while all other samples were diluted relative to this sample according to their β -actin/GAPDH average CT values.

Normalized RNA (5 μ l) was converted to cDNA and analyzed via TAQMAN® using One Step RT-PCR Master Mix Reagents (PE Biosystems; Catalog No. 4309169) and gene-specific primers according to the manufacturer's instructions. Probes and primers were designed for each assay according to Perkin Elmer Biosystem's *Primer Express* Software package (version I for Apple Computer's Macintosh Power PC) or a similar algorithm using the target sequence as input. Default settings were used for reaction conditions and the following parameters were set before selecting primers: primer concentration = 250 nM, primer melting temperature (T_m) range = 58°-60° C, primer optimal T_m = 59° C, maximum primer difference = 2° C, probe does not have 5' G, probe T_m must be 10° C greater than primer T_m , amplicon size 75 bp to 100 bp. The probes and primers selected (see below) were synthesized by SyntheGen (Houston, TX, USA). Probes were double purified by HPLC to remove uncoupled dye and evaluated by mass spectroscopy to verify coupling of reporter and quencher dyes to the 5' and 3' ends of the probe, respectively. Their final concentrations were: forward and reverse primers, 900 nM each, and probe, 200nM. The primers and probe used are designated Probe Name Ag1526 and are shown in Table 7.

Table 7. Probe Name: Ag1526

Primers	Sequences	T_m , °C	Length
Forward	5'-CTTGATGAGGGTCTCAATGG-3' (SEQ ID NO:7)	58.1	20
Probe	FAM-5'-CCACATCACTGAATTCCTGAGCATCC-3'-TAMRA (SEQ ID NO:8)	68.9	26
Reverse	5'-CAGACACAGTGAGCACCATG-3' (SEQ ID NO:9)	58.8	20

PCR conditions: Normalized RNA from each tissue and each cell line was spotted in each well of a 96 well PCR plate (Perkin Elmer Biosystems). PCR cocktails including two probes (SEQX-specific and another gene-specific probe multiplexed with the SEQX probe) were set up using 1X TaqMan™ PCR Master Mix for the PE Biosystems 7700, with 5 mM $MgCl_2$, dNTPs (dA, G, C, U at 1:1:1:2 ratios), 0.25 U/ml AmpliTaq Gold™ (PE Biosystems), and 0.4 U/ μ l RNase inhibitor, and 0.25 U/ μ l reverse transcriptase. Reverse transcription was performed at 48° C for 30 minutes followed by amplification/PCR cycles as follows: 95° C 10 min, then 40 cycles of 95° C for 15 seconds, 60° C for 1 minute.

The expression results are shown in Table 8.

Table 8.

Tissue_Name	Rel. Expr., %	Tissue_Name	Rel. Expr.,%
Endothelial cells	0.0	Renal carcinoma 786-0	0.1
Endothelial cells (treated)	0.5	Renal carcinoma A498	0.1
Pancreas	0.1	Renal carcinoma RXF 393	0.0
Pancreatic carcinoma CAPAN	13.9	Renal carcinoma ACHN	0.0
Adrenal Gland (new lot*)	0.0	Renal carcinoma UO-31	0.0
Thyroid	0.8	Renal carcinoma TK-10	0.3
Salivary gland	5.8	Liver	0.5
Pituitary gland	0.0	Liver (fetal)	1.9
Brain (fetal)	0.0	Liver carcinoma (hepatoblast) HepG2	22.5
Brain (whole)	0.0	Lung	0.4
Brain (amygdala)	0.0	Lung (fetal)	1.2
Brain (cerebellum)	0.0	Lung carcinoma (small cell) LX-1	12.2
Brain (hippocampus)	0.0	Lung carcinoma (small cell) NCI-H69	2.8
Brain (thalamus)	0.0	Lung carcinoma (small cell variant) SHP-77	0.0
Cerebral Cortex	0.1	Lung carcinoma (large cell) NCI-H460	0.0
Spinal cord	0.3	Lung carcinoma (non-small cell) A549	0.1
CNS carcinoma (glioma/astrocytoma) U87-MG	0.0	Lung carcinoma (non-small cell) NCI-H23	0.0
CNS carcinoma (glioma/astrocytoma) U-118-MG	0.0	Lung ca (non-small cell) HOP-62	0.0
CNS carcinoma (astrocytoma) SW1783	0.0	Lung carcinoma (non-small cell) NCI-H522	0.0
CNS carcinoma* (neuroblastoma; metastasis) SK-N-AS	0.0	Lung carcinoma (squamous) SW 900	10.6
CNS carcinoma (astrocytoma) SF-539	0.0	Lung carcinoma (squamous) NCI-H596	1.4
CNS carcinoma (astrocytoma) SNB-75	0.0	Mammary gland	0.5
CNS carcinoma (glioma) SNB-19	0.0	Breast carcinoma* (pleural effusion) MCF-7	63.1
CNS carcinoma (glioma) U251	0.0	Breast carcinoma* (pleural effusion) MDA-MB-231	0.0
CNS carcinoma (glioma) SF-29	0.0	Breast carcinoma* (pleural effusion) T47D	0.6
Heart	1.6	Breast carcinoma BT-549	0.0
Skeletal Muscle (new lot*)	0.0	Breast carcinoma MDA-N	0.0
Bone marrow	0.0	Ovary	0.2
Thymus	1.9	Ovarian carcinoma OVCAR-3	2.8
Spleen	0.0	Ovarian carcinoma OVCAR-4	79.9
Lymph node	0.0	Ovarian carcinoma OVCAR-5	5.3
Colorectal	3.6	Ovarian carcinoma OVCAR-8	0.1
Stomach	1.5	Ovarian carcinoma IGROV-1	2.1

Small intestine	3.5	Ovarian carcinoma* (ascites) SK-OV-3	0.9
Colon carcinoma SW480	0.3	Uterus	0.2
Colon carcinoma* (SW480 metastasis)SW620	4.3	Placenta	0.0
Colon carcinoma HT29	12.0	Prostate	1.8
Colon carcinoma HCT-116	31.4	Prostate carcinoma* (bone metastasis)PC-3	0.4
Colon carcinoma CaCo-2	44.2	Testis	0.0
83219 CC Well to Mod Diff (ODO3866)	9.8	Melanoma Hs688(A).T	0.0
Colon carcinoma HCC-2998	100.0	Melanoma* (metastasis) Hs688(B).T	0.0
Gastric carcinoma* (liver metastasis) NCI-N87	37.8	Melanoma UACC-62	0.0
Bladder	2.2	Melanoma M14	0.0
Trachea	0.4	Melanoma LOX IMVI	0.0
Kidney	4.6	Melanoma* (metastasis) SK-MEL-5	0.0
Kidney (fetal)	0.7	Renal carcinoma 786-0	0.1

* = established from metastasis

The results in Table 8 indicate that the clone of SEQ ID NO:5 is very strongly expressed in several tumor derived cell lines compared with normal tissue, especially colon tumor cells, breast tumor cells and ovarian tumor cells. This observation indicates that the clone of SEQ ID NO:5 has a role in cell proliferation and potential utility as a blood marker to identify and/or stage tumors. It may furthermore be a target for a specific monoclonal antibody that could be used to treat various cancers, especially colon cancer, breast cancer and ovarian cancer.

Example 4. Real Time Quantitative Expression of Clone 65677221-3-frag in Surgical Tumor Tissues

Real time quantitative expression of Clone 65677221-3-frag was examined in a 96 well (2 control wells, 94 test samples) panel of RNA or cDNA samples isolated from human tissues. The tissues were derived from human malignancies; in cases where indicated many malignant tissues are accompanied by "matched margins". These matched margins were taken from the tissue surrounding (i.e. immediately proximal) to the zone of surgery (designated "NAT", for normal adjacent tissue, in Table 9). The tumor tissue and the "matched margins" are evaluated by two independent pathologists (the surgical pathologists and again by a pathologists at NDRI or CHTN). This analysis provides a gross histopathological assessment of tumor differentiation grade. Moreover, most samples include

the original surgical pathology report that provides information regarding the clinical stage of the patient. Additional RNA or cDNA samples were obtained from various human tissues derived from autopsies performed on elderly people or sudden death human victims. These tissue were ascertained to be free of disease and were purchased from various commercial
5 sources such as Clontech (Palo Alto, CA), Research Genetics, and Invitrogen (Carlsbad CA).

RNA integrity from all samples was determined by visual assessment of agarose gel electrophoresis using 28s and 18s ribosomal RNA staining intensity ratio as a guide (2:1 to 2.5:1 28s:18s). Integrity was also determined by verifying the absence of low molecular weight RNAs, which is indicative of degradation products. Samples were examined for
10 genomic DNA contamination by reactions performed in the absence of reverse transcriptase using probe and primer sets designed to amplify across the span of a single exon.

The same primer-probe set, Ag1526, was used as in Example 3. The results are presented in Table 9.

Table 9

Tissue_Name	Rel. Expr., %	Tissue_Name	Rel. Expr., %
Normal Colon GENPAK 061003	22.4	Kidney NAT Clontech 8120608	6.4
83219 CC Well to Mod Diff (ODO3866)	32.0	Kidney Cancer Clontech 8120613	0.3
83220 CC NAT (ODO3866)	11.5	Kidney NAT Clontech 8120614	7.6
83221 CC Gr.2 rectosigmoid (ODO3868)	19.1	Kidney Cancer Clontech 9010320	0.1
83222 CC NAT (ODO3868)	2.7	Kidney NAT Clontech 9010321	3.9
83235 CC Mod Diff (ODO3920)	5.7	Normal Uterus GENPAK 061018	0.0
83236 CC NAT (ODO3920)	5.2	Uterus Cancer GENPAK 064011	0.0
83237 CC Gr.2 ascend colon (ODO3921)	6.9	Normal Thyroid Clontech A+ 6570-1**	10.1
83238 CC NAT (ODO3921)	10.8	Thyroid Cancer GENPAK 064010	79.8
83241 CC from Partial Hepatectomy (ODO4309)	9.7	Thyroid Cancer INVITROGEN A302152	13.3
83242 Liver NAT (ODO4309)	0.5	Thyroid NAT INVITROGEN A302153	5.9
87472 Colon mets to lung (OD04451-01)	28.2	Normal Breast GENPAK 061019	5.5
87473 Lung NAT (OD04451-02)	24.5	84877 Breast Cancer (OD04566)	36.5
Normal Prostate Clontech A+ 6546-1	3.3	85975 Breast Cancer (OD04590-01)	9.7
84140 Prostate Cancer (OD04410)	0.1	85976 Breast Cancer Mets (OD04590-03)	7.3
84141 Prostate NAT (OD04410)	0.1	87070 Breast Cancer Metastasis (OD04655-05)	2.3
87073 Prostate Cancer (OD04720-01)	0.2	GENPAK Breast Cancer 064006	7.4
87074 Prostate NAT (OD04720-02)	0.3	Breast Cancer Clontech 9100266	66.7
Normal Lung GENPAK 061010	8.4	Breast NAT Clontech 9100265	39.8
83239 Lung Met to Muscle (ODO4286)	1.8	Breast Cancer INVITROGEN A209073	15.4
83240 Muscle NAT (ODO4286)	0.0	Breast NAT INVITROGEN A2090734	7.6
84136 Lung Malignant Cancer (OD03126)	6.7	Normal Liver GENPAK 061009	3.5
84137 Lung NAT (OD03126)	0.6	Liver Cancer GENPAK 064003	0.3
84871 Lung Cancer (OD04404)	12.0	Liver Cancer Research Genetics RNA 1025	6.7
84872 Lung NAT (OD04404)	7.1	Liver Cancer Research Genetics RNA 1026	3.4
84875 Lung Cancer (OD04565)	28.2	Paired Liver Cancer Tissue Research Genetics RNA 6004-T	7.0
85950 Lung Cancer (OD04237-	0.8	Paired Liver Tissue Research	4.9

01)		Genetics RNA 6004-N	
85970 Lung NAT (OD04237-02)	0.9	Paired Liver Cancer Tissue Research Genetics RNA 6005-T	4.3
83255 Ocular Mel Met to Liver (ODO4310)	0.0	Paired Liver Tissue Research Genetics RNA 6005-N	5.3
83256 Liver NAT (ODO4310)	0.3	Normal Bladder GENPAK 061001	16.3
84139 Melanoma Mets to Lung (OD04321)	0.1	Bladder Cancer Research Genetics RNA 1023	32.5
84138 Lung NAT (OD04321)	4.2	Bladder Cancer INVITROGEN A302173	44.5
Normal Kidney GENPAK 061008	2.9	87071 Bladder Cancer (OD04718-01)	2.3
83786 Kidney Ca, Nuclear grade 2 (OD04338)	0.2	87072 Bladder Normal Adjacent (OD04718-03)	0.0
83787 Kidney NAT (OD04338)	1.6	Normal Ovary Res. Gen.	1.0
83788 Kidney Ca Nuclear grade 1/2 (OD04339)	0.3	Ovarian Cancer GENPAK 064008	9.6
83789 Kidney NAT (OD04339)	0.5	87492 Ovary Cancer (OD04768-07)	9.1
83790 Kidney Ca, Clear cell type (OD04340)	1.2	87493 Ovary NAT (OD04768-08)	0.0
83791 Kidney NAT (OD04340)	1.7	Normal Stomach GENPAK 061017	4.6
83792 Kidney Ca, Nuclear grade 3 (OD04348)	1.7	NAT Stomach Clontech 9060359	41.0
83793 Kidney NAT (OD04348)	0.6	Gastric Cancer Clontech 9060395	15.1
87474 Kidney Cancer (OD04622-01)	0.0	NAT Stomach Clontech 9060394	48.7
87475 Kidney NAT (OD04622-03)	5.3	Gastric Cancer Clontech 9060397	78.6
85973 Kidney Cancer (OD04450-01)	2.4	NAT Stomach Clontech 9060396	100.0
85974 Kidney NAT (OD04450-03)	1.7	Gastric Cancer GENPAK 064005	15.2
Kidney Cancer Clontech 8120607	24.5	Kidney NAT Clontech 8120608	6.4

- The results shown in Table 9 demonstrate that clone 65677221-3-frag is strongly upregulated by cancer cell lines and tumor, compared to normal tissue and normal adjacent tissues. This is especially true for cancers of the breast, ovary, colon, stomach, and pancreas.
- 5 Accordingly, growth, migration and/or invasion of tumors for which FCTR_X nucleic acids are upregulated (see, *e.g.*, Table 9) can be inhibited by antagonizing expression or activity of a FCTR_X nucleic acid or polypeptide, respectively. For example, a monoclonal antibody to a FCTR_X polypeptide can be used to inhibit tumor cell growth, cell migration and/or invasion.

Example 5. Real Time Quantitative Expression of Clone 65677221-3-frag in Tissues Exposed to an Inflammatory Agent

Real Time Quantitative Expression of Clone 65677221-3-frag in was examined in a 96 well plate (2 control wells, 94 test samples) composed of RNA or cDNA isolated from various human cell lines or tissues exposed to inflammatory agents. Total RNA from normal tissues (colon and lung, purchased from Stratagene (La Jolla, CA), and thymus and kidney purchased from Clontech (Palo Alto, CA)) was used as controls. Total RNA from liver tissue from cirrhosis patients and kidney RNA from Lupus patients were obtained from Biochain (Hayward, CA). Intestinal tissue for RNA preparation from tissues Crohns disease and ulcerative colitis patients was obtained from the National Disease Research Interchange (NDRI) (Philadelphia, PA).

Astrocytes, lung fibroblasts, dermal fibroblasts, coronary artery smooth muscle cells, small airway epithelium, bronchial epithelium, microvascular dermal endothelial cells, microvascular lung endothelial cells, human pulmonary aortic endothelial cells, human umbilical vein endothelial cells were all purchased from Clonetics (Walkersville, MD) and grown in the media supplied for these cell types by Clonetics. These primary cell types were activated with various cytokines or combinations of cytokines for 6 and/or 12-14 hours. The following cytokines were used; IL-1 beta at approximately 1-5 ng/ml, TNF alpha at approximately 5-10 ng/ml, IFN gamma at approximately 20-50 ng/ml, IL-4 at approximately 5-10 ng/ml, IL-9 at approximately 5-10 ng/ml, IL-13 at approximately 5-10 ng/ml. For endothelial cells sometimes were starved for various times by culture in the basal media from Clonetics with 0.1% serum.

Mononuclear cells were prepared from blood of human subjects using Ficoll. LAK cells were prepared from these cells by culture in DMEM 5% FCS (Hyclone), 100 μ M non essential amino acids (Gibco/Life Technologies, Rockville, MD), 1 mM sodium pyruvate (Gibco), mercaptoethanol 5.5×10^{-5} M (Gibco), and 10 mM Hepes (Gibco) and Interleukin 2 for 4-6 days. Cells were then either activated with 10-20 ng/ml PMA and 1-2 μ g/ml ionomycin, IL-12 at 5-10 ng/ml, IFN gamma at 20-50 ng/ml and IL-18 at 5-10 ng/ml for 6 hours. In some cases, mononuclear cells were cultured for 4-5 days in DMEM 5% FCS (Hyclone), 100 μ M non essential amino acids (Gibco), 1 mM sodium pyruvate (Gibco), mercaptoethanol 5.5×10^{-5} M (Gibco), and 10 mM Hepes (Gibco) with PHA or PWM at approximately 5 μ g/ml. Samples were taken at 24, 48 and 72 hours for RNA preparation. MLR samples were obtained by taking blood from two donors, isolating the mononuclear cells using Ficoll and mixing the isolated mononuclear cells 1:1 at a final concentration of

approximately 2×10^6 cells/ml in DMEM 5% FCS (Hyclone), 100 μ M non essential amino acids (Gibco), 1 mM sodium pyruvate (Gibco), mercaptoethanol (5.5×10^{-5} M) (Gibco), and 10 mM Hepes (Gibco). The MLR was cultured and samples taken at various time points ranging from 1- 7 days for RNA preparation.

5 To prepare monocytes, macrophages and dendritic cells, monocytes were isolated from mononuclear cells using CD14 Miltenyi Beads, positive VS selection columns and a Vario Magnet as per the manufacturer's instructions. Monocytes were differentiated into dendritic cells by culture in DMEM 5% FCS (Hyclone, Logan, UT), 100 μ M non essential amino acids (Gibco), 1 mM sodium pyruvate (Gibco), mercaptoethanol 5.5×10^{-5} M (Gibco), and 10 mM
10 Hepes (Gibco), 50 ng/ml GMCSF and 5 ng/ml IL-4 for 5-7 days. Macrophages were prepared by culture of monocytes for 5-7 days in DMEM 5% FCS (Hyclone), 100 μ M non essential amino acids (Gibco), 1 mM sodium pyruvate (Gibco), mercaptoethanol 5.5×10^{-5} M (Gibco), 10 mM Hepes (Gibco) and 10% AB Human Serum or MCSF at approximately 50 ng/ml. Monocytes, macrophages and dendritic cells were stimulated for 6 and 12-14 hours with LPS
15 at 100 ng/ml. Dendritic cells were also stimulated with anti-CD40 monoclonal antibody (Pharmingen) at 10 μ g/ml for 6 and 12-14 hours.

CD4 lymphocytes, CD8 lymphocytes and NK cells were also isolated from mononuclear cells using CD4, CD8 and CD56 Miltenyi beads, positive VS selection columns and a Vario Magnet as per the manufacturer's instructions. CD45RA and CD45RO CD4
20 lymphocytes were isolated by depleting mononuclear cells of CD8, CD56, CD14 and CD19 cells using CD8, CD56, CD14 and CD19 Miltenyi beads and +ve selection. Then CD4RO beads were used to isolate the CD45RO CD4 lymphocytes with the remaining cells being CD45RA CD4 lymphocytes. CD45RA CD4, CD45RO CD4 and CD8 lymphocytes were placed in DMEM 5% FCS (Hyclone), 100 μ M non essential amino acids (Gibco), 1 mM
25 sodium pyruvate (Gibco), mercaptoethanol 5.5×10^{-5} M (Gibco), and 10 mM Hepes (Gibco) and plated at 10^6 cells/ml onto Falcon 6 well tissue culture plates that had been coated overnight with 0.5 μ g/ml anti-CD28 (Pharmingen) and 3 μ g/ml anti-CD3 (OKT3, ATCC) in PBS. After 6 and 24 hours, the cells were harvested for RNA preparation. To prepare chronically activated CD8 lymphocytes, we activated the isolated CD8 lymphocytes for 4 days
30 on anti-CD28 and anti-CD3 coated plates and then harvested the cells and expanded them in DMEM 5% FCS (Hyclone), 100 μ M non essential amino acids (Gibco), 1 mM sodium pyruvate (Gibco), mercaptoethanol 5.5×10^{-5} M (Gibco), and 10 mM Hepes (Gibco) and IL-2. The expanded CD8 cells were then activated again with plate bound anti-CD3 and anti-CD28 for 4 days and expanded as before. RNA was isolated 6 and 24 hours after the second

activation and after 4 days of the second expansion culture. The isolated NK cells were cultured in DMEM 5% FCS (Hyclone), 100 μ M non essential amino acids (Gibco), 1 mM sodium pyruvate (Gibco), mercaptoethanol 5.5×10^{-5} M (Gibco), and 10 mM Hepes (Gibco) and IL-2 for 4-6 days before RNA was prepared.

5 B cells were obtained from tonsils. Tonsils were cut with sterile dissecting scissors and then passed through a sieve. Tonsil cells were then pelleted and resuspended at 10^6 cells/ml in DMEM 5% FCS (Hyclone), 100 μ M non essential amino acids (Gibco), 1 mM sodium pyruvate (Gibco), mercaptoethanol 5.5×10^{-5} M (Gibco), and 10 mM Hepes (Gibco). Cells were activated with PWM at 5 μ g/ml or anti-CD40 (Pharmingen) at approximately 10
10 μ g/ml and IL-4 at 5-10 ng/ml. Cells were harvested for RNA preparation at 24, 48 and 72 hours.

To prepare the primary and secondary Th1/Th2 and Tr1 cells, six-well Falcon plates were coated overnight with 10 μ g/ml anti-CD28 (Pharmingen) and 2 μ g/ml OKT3 (ATCC), and then washed twice with PBS. Umbilical cord blood CD4 lymphocytes (Poietic Systems,
15 German Town, MD) were cultured at 10^5 - 10^6 cells/ml in DMEM 5% FCS (Hyclone), 100 μ M non essential amino acids (Gibco), 1 mM sodium pyruvate (Gibco), mercaptoethanol 5.5×10^{-5} M (Gibco), 10 mM Hepes (Gibco) and IL-2 (4 ng/ml). IL-12 (5 ng/ml) and anti-IL4 (1 g/ml) were used to direct to Th1, while IL-4 (5 ng/ml) and anti-IFN gamma (1 g/ml) were used to direct to Th2 and IL-10 at 5 ng/ml was used to direct to Tr1. After 4-5 days, the
20 activated Th1, Th2 and Tr1 lymphocytes were washed once in DMEM and expanded for 4-7 days in DMEM 5% FCS (Hyclone), 100 μ M non essential amino acids (Gibco), 1 mM sodium pyruvate (Gibco), mercaptoethanol 5.5×10^{-5} M (Gibco), 10 mM Hepes (Gibco) and IL-2 (1 ng/ml). Following this, the activated Th1, Th2 and Tr1 lymphocytes were re-stimulated for 5 days with anti-CD28/OKT3 and cytokines as described above, but with the addition of anti-
25 CD95L (1 g/ml) to prevent apoptosis. After 4-5 days, the Th1, Th2 and Tr1 lymphocytes were washed and then expanded again with IL-2 for 4-7 days. Activated Th1 and Th2 lymphocytes were maintained in this way for a maximum of three cycles. RNA was prepared from primary and secondary Th1, Th2 and Tr1 after 6 and 24 hours following the second and third activations with plate bound anti-CD3 and anti-CD28 mAbs and 4 days into the second
30 and third expansion cultures in Interleukin 2.

The following leukocyte cell lines were obtained from the ATCC (Manassas, VA): Ramos, EOL-1, KU-812. EOL cells were further differentiated by culture in 0.1 mM dbcAMP at 5×10^5 cells/ml for 8 days, changing the media every 3 days and adjusting the cell concentration to 5×10^5 cells/ml. The cells were cultured with DMEM or RPMI (as

recommended by the ATCC), with the addition of 5% FCS (Hyclone), 100 μ M non essential amino acids (Gibco), 1 mM sodium pyruvate (Gibco), mercaptoethanol 5.5×10^{-5} M (Gibco), 10 mM Hepes (Gibco). RNA was either prepared from resting cells or cells activated with PMA at 10 ng/ml and ionomycin at 1 μ g/ml for 6 and 14 hours. A keratinocyte line CCD106 and an airway epithelial tumor line NCI-H292 were also obtained from the ATCC. Both were cultured in DMEM 5% FCS (Hyclone), 100 μ M non essential amino acids (Gibco), 1 mM sodium pyruvate (Gibco), mercaptoethanol 5.5×10^{-5} M (Gibco), and 10 mM Hepes (Gibco). CCD1106 cells were activated for 6 and 14 hours with approximately 5 ng/ml TNF alpha and 1 ng/ml IL-1 beta, while NCI-H292 cells were activated for 6 and 14 hours with the following cytokines: 5 ng/ml IL-4, 5 ng/ml IL-9, 5 ng/ml IL-13 and 25 ng/ml IFN gamma.

For these cell lines and blood cells, RNA was prepared by lysing approximately 10^7 cells/ml using Trizol (Gibco). Briefly, 1/10 volume of bromochloropropane (Molecular Research Corporation) was added to the RNA sample, vortexed and after 10 minutes at room temperature, the tubes were spun at 14,000 rpm in a Sorvall SS34 rotor. The aqueous phase was removed and placed in a 15 ml Falcon Tube. An equal volume of isopropanol was added and left at -20 degrees C overnight. The precipitated RNA was pelleted at 9,000 rpm for 15 min in a Sorvall SS34 rotor and washed in 70% ethanol. The pellet was redissolved in 300 μ l of RNase-free water and 35 μ l buffer (Promega, Madison, WI) 5 μ l DTT, 7 μ l RNasin and 8 μ l DNase were added. The tube was incubated at 37 degrees C for 30 minutes to remove contaminating genomic DNA, extracted once with phenol chloroform and re-precipitated with 1/10 volume of 3 M sodium acetate and 2 volumes of 100% ethanol. The RNA was spun down and placed in RNase free water. RNA was stored at -80 degrees C.

The same primer-probe set, Ag1526, was used as in Example 3. The results are presented in Table 10.

Table 10.

Tissue_Name	Rel. Expr., %	Tissue_Name	Rel. Expr., %
93768_Secondary Th1_anti-CD28/anti-CD3	0.0	93100_HUVEC (Endothelial)_IL-1b	0.0
93769_Secondary Th2_anti-CD28/anti-CD3	0.0	93779_HUVEC (Endothelial)_IFN gamma	0.0
93770_Secondary Tr1_anti-CD28/anti-CD3	0.0	93102_HUVEC (Endothelial)_TNF alpha + IFN gamma	0.0
93573_Secondary Th1_resting day 4-6 in IL-2	0.0	93101_HUVEC (Endothelial)_TNF alpha + IL4	0.0
93572_Secondary Th2_resting day 4-6	0.0	93781_HUVEC (Endothelial)_IL-11	0.0

in IL-2			
93571_Secondary Tr1_resting day 4-6 in IL-2	0.0	93583_Lung Microvascular Endothelial Cells_none	0.0
93568_primary Th1_anti-CD28/anti-CD3	0.0	93584_Lung Microvascular Endothelial Cells_TNFa (4 ng/ml) and IL1b (1 ng/ml)	0.0
93569_primary Th2_anti-CD28/anti-CD3	0.0	92662_Microvascular Dermal endothelium_none	0.0
93570_primary Tr1_anti-CD28/anti-CD3	0.0	92663_Microvascular Dermal endothelium_TNFa (4 ng/ml) and IL1b (1 ng/ml)	0.0
93565_primary Th1_resting dy 4-6 in IL-2	0.0	93773_Bronchial epithelium_TNFa (4 ng/ml) and IL1b (1 ng/ml) **	66.0
93566_primary Th2_resting dy 4-6 in IL-2	0.0	93347_Small Airway Epithelium_none	29.9
93567_primary Tr1_resting dy 4-6 in IL-2	0.0	93348_Small Airway Epithelium_TNFa (4 ng/ml) and IL1b (1 ng/ml)	100.0
93351_CD45RA CD4 lymphocyte_anti-CD28/anti-CD3	0.0	92668_Coronary Artery SMC_resting	0.0
93352_CD45RO CD4 lymphocyte_anti-CD28/anti-CD3	0.0	92669_Coronary Artery SMC_TNFa (4 ng/ml) and IL1b (1 ng/ml)	0.0
93251_CD8 Lymphocytes_anti-CD28/anti-CD3	0.0	93107_astrocytes_resting	0.0
93353_chronic CD8 Lymphocytes 2ry_resting dy 4-6 in IL-2	0.0	93108_astrocytes_TNFa (4 ng/ml) and IL1b (1 ng/ml)	0.0
93574_chronic CD8 Lymphocytes 2ry_activated CD3/CD28	0.0	92666_KU-812 (Basophil)_resting	0.0
93354_CD4_none	0.0	92667_KU-812 (Basophil)_PMA/ionomycin	0.0
93252_Secondary Th1/Th2/Tr1_anti-CD95 CH11	0.0	93579_CCD1106 (Keratinocytes)_none	35.9
93103_LAK cells_resting	0.0	93580_CCD1106 (Keratinocytes)_TNFa and IFNg	86.5
93788_LAK cells_IL-2	0.0	93791_Liver Cirrhosis	0.4
93787_LAK cells_IL-2+IL-12	0.0	93792_Lupus Kidney	0.4
93789_LAK cells_IL-2+IFN gamma	0.0	93577_NCI-H292	9.3
93790_LAK cells_IL-2+ IL-18	0.0	93358_NCI-H292_IL-4	8.0
93104_LAK cells_PMA/ionomycin and IL-18	0.0	93360_NCI-H292_IL-9	9.1
93578_NK Cells IL-2_resting	0.0	93359_NCI-H292_IL-13	6.9
93109_Mixed Lymphocyte Reaction_Two Way MLR	0.0	93357_NCI-H292_IFN gamma	5.2
93110_Mixed Lymphocyte Reaction_Two Way MLR	0.0	93777_HPAEC_-	0.0
93111_Mixed Lymphocyte Reaction_Two Way MLR	0.0	93778_HPAEC_IL-1 beta/TNA alpha	0.0
93112_Mononuclear Cells (PBMCs)_resting	0.0	93254_Normal Human Lung Fibroblast_none	0.0

93113_Mononuclear Cells (PBMcs)_PWM	0.0	93253_Normal Human Lung Fibroblast_TNFa (4 ng/ml) and IL-1b (1 ng/ml)	0.0
93114_Mononuclear Cells (PBMcs)_PHA-L	0.0	93257_Normal Human Lung Fibroblast_IL-4	0.0
93249_Ramos (B cell)_none	0.0	93256_Normal Human Lung Fibroblast_IL-9	0.0
93250_Ramos (B cell)_ionomycin	0.0	93255_Normal Human Lung Fibroblast_IL-13	0.0
93349_B lymphocytes_PWM	0.0	93258_Normal Human Lung Fibroblast_IFN gamma	0.1
93350_B lymphocytes_CD40L and IL-4	0.0	93106_Dermal Fibroblasts CCD1070_resting	0.0
92665_EOL-1 (Eosinophil)_dbcAMP differentiated	0.0	93361_Dermal Fibroblasts CCD1070_TNF alpha 4 ng/ml	0.0
93248_EOL-1 (Eosinophil)_dbcAMP/PmAionomycin	0.0	93105_Dermal Fibroblasts CCD1070_IL-1 beta 1 ng/ml	0.0
93356_Dendritic Cells_none	0.0	93772_dermal fibroblast_IFN gamma	0.0
93355_Dendritic Cells_LPS 100 ng/ml	0.0	93771_dermal fibroblast_IL-4	0.0
93775_Dendritic Cells_anti-CD40	0.0	93259_IBD Colitis 1**	1.7
93774_Monocytes_resting	0.0	93260_IBD Colitis 2	0.0
93776_Monocytes_LPS 50 ng/ml	0.0	93261_IBD Crohns	0.2
93581_Macrophages_resting	0.0	735010_Colon_normal	4.8
93582_Macrophages_LPS 100 ng/ml	0.0	735019_Lung_none	1.4
93098_HUVEC (Endothelial)_none	0.0	64028-1_Thymus_none	0.7
93099_HUVEC (Endothelial)_starved	0.0	64030-1_Kidney_none	4.5

5 The results shown in Table 10 demonstrate that expression of clone 65677221-3-frag is increased by TNF-alpha in epithelial airways. Accordingly, inflammatory conditions can be treated or prevented by antagonizing expression or activity of a FCTR_X nucleic acid or polypeptide, respectively. For example, inflammatory conditions in the lung such as asthma, as well as other lung-inflammation diseases, can be prevented or treated by inhibiting expression or activity of a FCTR_X nucleic acid or polypeptide.

10

OTHER EMBODIMENTS

Although particular embodiments have been disclosed herein in detail, this has been done by way of example for purposes of illustration only, and is not intended to be limiting with respect to the scope of the appended claims. In particular, it is contemplated by the inventor that various substitutions, alterations, and modifications may be made to the invention without departing from the spirit and scope of the invention as defined by the claims. Therefore, other aspects, advantages, and modifications are within the scope of the following claims.